

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| Effectiveness of Care: Prevention and Screening | | | | | | |
| Adult BMI Assessment (aba) | Y | | N | 72.44% | R | |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (wcc) | Y | | N | | | |
| <i>BMI Percentile</i> | | | | 46.99% | R | |
| <i>Counseling for Nutrition</i> | | | | 41.43% | R | |
| <i>Counseling for Physical Activity</i> | | | | 26.95% | R | |
| Childhood Immunization Status (cis) | Y | | N | | | |
| <i>DTaP</i> | | | | 83.22% | R | |
| <i>IPV</i> | | | | 94.04% | R | |
| <i>MMR</i> | | | | 92.49% | R | |
| <i>HIB</i> | | | | 93.82% | R | |
| <i>Hepatitis B</i> | | | | 91.39% | R | |
| <i>VZV</i> | | | | 91.39% | R | |
| <i>Pneumococcal Conjugate</i> | | | | 84.11% | R | |
| <i>Hepatitis A</i> | | | | 89.40% | R | |
| <i>Rotavirus</i> | | | | 75.28% | R | |
| <i>Influenza</i> | | | | 53.64% | R | |
| <i>Combination #2</i> | | | | 79.47% | R | |
| <i>Combination #3</i> | | | | 76.60% | R | |
| <i>Combination #4</i> | | | | 74.61% | R | |
| <i>Combination #5</i> | | | | 65.34% | R | |
| <i>Combination #6</i> | | | | 46.80% | R | |
| <i>Combination #7</i> | | | | 63.80% | R | |
| <i>Combination #8</i> | | | | 45.70% | R | |
| <i>Combination #9</i> | | | | 41.50% | R | |
| <i>Combination #10</i> | | | | 40.62% | R | |
| Immunizations for Adolescents (ima) | Y | | | | | |
| <i>Meningococcal</i> | | | | NR | NR | |
| <i>Tdap/Td</i> | | | | NR | NR | |
| <i>Combination #1</i> | | | | NR | NR | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| Human Papillomavirus Vaccine for Female Adolescents (hpv) | Y | | | 9.17% | R | |
| Lead Screening in Children (lsc) | Y | | N | 34.07% | R | |
| Breast Cancer Screening (bcs) | Y | | | 55.60% | R | |
| Cervical Cancer Screening (ccs) | Y | | | 58.44% | R | |
| Chlamydia Screening in Women (chl) | Y | | | | | |
| 16-20 Years | | | | 40.39% | R | |
| 21-24 Years | | | | 54.45% | R | |
| Total | | | | 45.87% | R | |
| Effectiveness of Care: Respiratory Conditions | | | | | | |
| Appropriate Testing for Children with Pharyngitis (cwp) | Y | Y | | 63.48% | R | |
| Appropriate Treatment for Children With URI (uri) | Y | Y | | 84.96% | R | |
| Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (aab) | Y | Y | | 17.58% | R | |
| Use of Spirometry Testing in the Assessment and Diagnosis of COPD (spr) | Y | | | NA | R | |
| Pharmacotherapy Management of COPD Exacerbation (pce) | Y | Y | | | | |
| Systemic Corticosteroid | | | | NA | R | |
| Bronchodilator | | | | NA | R | |
| Use of Appropriate Medications for People With Asthma (asm) | Y | Y | | | | |
| 5-11 Years | | | | 93.90% | R | |
| 12-18 Years | | | | 77.78% | R | |
| 19-50 Years | | | | 86.05% | R | |
| 51-64 Years | | | | NA | R | |
| Total | | | | 86.46% | R | |
| Medication Management for People With Asthma (mma) | Y | Y | | | | |
| 5-11 Years - Medication Compliance 50% | | | | 44.16% | R | |
| 5-11 Years - Medication Compliance 75% | | | | 23.38% | R | |
| 12-18 Years - Medication Compliance 50% | | | | 32.65% | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| 12-18 Years - Medication Compliance 75% | | | | 16.33% | R | |
| 19-50 Years - Medication Compliance 50% | | | | 45.95% | R | |
| 19-50 Years - Medication Compliance 75% | | | | 29.73% | R | |
| 51-64 Years - Medication Compliance 50% | | | | NA | R | |
| 51-64 Years - Medication Compliance 75% | | | | NA | R | |
| Total - Medication Compliance 50% | | | | 42.17% | R | |
| Total - Medication Compliance 75% | | | | 22.89% | R | |
| Asthma Medication Ratio (amr) | Y | | | | | |
| 5-11 Years | | | | 65.85% | R | |
| 12-18 Years | | | | 41.27% | R | |
| 19-50 Years | | | | 48.84% | R | |
| 51-64 Years | | | | NA | R | |
| Total | | | | 54.17% | R | |
| Effectiveness of Care: Cardiovascular | | | | | | |
| Cholesterol Management for Patients With Cardiovascular Conditions (cmc) | Y | | N | | | |
| LDL-C Screening Performed | | | | NA | R | |
| LDL-C Control (<100 mg/dL) | | | | NA | R | |
| Controlling High Blood Pressure (cbp) | Y | | | 49.06% | R | |
| Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) | Y | Y | | NA | R | |
| Effectiveness of Care: Diabetes | | | | | | |
| Comprehensive Diabetes Care (cdc) | Y | | Y | | | |
| Hemoglobin A1c (HbA1c) Testing | | | | 82.33% | R | |
| HbA1c Poor Control (>9.0%) | | | | 39.66% | R | |
| HbA1c Control (<8.0%) | | | | 45.69% | R | |
| HbA1c Control (<7.0%) | | | | NR | NR | |
| Eye Exam (Retinal) Performed | | | | 53.88% | R | |
| LDL-C Screening Performed | | | | 72.41% | R | |
| LDL-C Control (<100 mg/dL) | | | | 39.22% | R | |
| Medical Attention for Nephropathy | | | | 76.29% | R | |
| Blood Pressure Control (<140/80 mm Hg) | | | | 46.55% | R | |
| Blood Pressure Control (<140/90 mm Hg) | | | | 65.09% | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| Effectiveness of Care: Musculoskeletal | | | | | | |
| Disease Modifying Anti-Rheumatic Drug Therapy in Rheumatoid Arthritis (art) | Y | Y | | NA | R | |
| Use of Imaging Studies for Low Back Pain (lbp) | Y | | | 79.86% | R | |
| Effectiveness of Care: Behavioral Health | | | | | | |
| Antidepressant Medication Management (amm) | N | N | | | | |
| <i>Effective Acute Phase Treatment</i> | | | | NR | NR | |
| <i>Effective Continuation Phase Treatment</i> | | | | NR | NR | |
| Follow-Up Care for Children Prescribed ADHD Medication (add) | Y | Y | | | | |
| <i>Initiation Phase</i> | | | | 44.74% | R | |
| <i>Continuation and Maintenance (C&M) Phase</i> | | | | NA | R | |
| Follow-Up After Hospitalization for Mental Illness (fuh) | N | N | | | | |
| <i>30-Day Follow-Up</i> | | | | NR | NR | |
| <i>7-Day Follow-Up</i> | | | | NR | NR | |
| Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (ssd) | Y | Y | | NA | R | |
| Diabetes Monitoring for People With Diabetes and Schizophrenia (smd) | Y | | | NA | R | |
| Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (smc) | Y | | | NA | R | |
| Adherence to Antipsychotic Medications for Individuals With Schizophrenia (saa) | Y | Y | | NA | R | |
| Effectiveness of Care: Medication Management | | | | | | |
| Annual Monitoring for Patients on Persistent Medications (mpm) | Y | Y | | | | |
| <i>ACE Inhibitors or ARBs</i> | | | | 89.32% | R | |
| <i>Digoxin</i> | | | | NA | R | |
| <i>Diuretics</i> | | | | 86.30% | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| <i>Anticonvulsants</i> | | | | 61.76% | R | |
| <i>Total</i> | | | | 86.12% | R | |
| Access/Availability of Care | | | | | | |
| Adults' Access to Preventive/Ambulatory Health Services (aap) | Y | | | | | |
| <i>20-44 Years</i> | | | | 81.70% | R | |
| <i>45-64 Years</i> | | | | 85.04% | R | |
| <i>65+ Years</i> | | | | NA | R | |
| <i>Total</i> | | | | 82.57% | R | |
| Children and Adolescents' Access to Primary Care Practitioners (cap) | Y | | | | | |
| <i>12-24 Months</i> | | | | 97.20% | R | |
| <i>25 Months - 6 Years</i> | | | | 86.12% | R | |
| <i>7-11 Years</i> | | | | 85.22% | R | |
| <i>12-19 Years</i> | | | | 85.76% | R | |
| Annual Dental Visit (adv) | Y | Y | | | | |
| <i>2-3 Years</i> | | | | 55.74% | R | |
| <i>4-6 Years</i> | | | | 68.53% | R | |
| <i>7-10 Years</i> | | | | 73.35% | R | |
| <i>11-14 Years</i> | | | | 66.18% | R | |
| <i>15-18 Years</i> | | | | 56.43% | R | |
| <i>19-21 Years</i> | | | | 37.45% | R | |
| <i>Total</i> | | | | 64.00% | R | |
| Initiation and Engagement of AOD Dependence Treatment (iet) | N | N | | | | |
| <i>Initiation of AOD Treatment: 13-17 Years</i> | | | | NR | NR | |
| <i>Engagement of AOD Treatment: 13-17 Years</i> | | | | NR | NR | |
| <i>Initiation of AOD Treatment: 18+ Years</i> | | | | NR | NR | |
| <i>Engagement of AOD Treatment: 18+ Years</i> | | | | NR | NR | |
| <i>Initiation of AOD Treatment: Total</i> | | | | NR | NR | |
| <i>Engagement of AOD Treatment: Total</i> | | | | NR | NR | |
| Prenatal and Postpartum Care (ppc) | Y | | | | | |
| <i>Timeliness of Prenatal Care</i> | | | | 86.06% | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|--------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| <i>Postpartum Care</i> | | | | 63.05% | R | |
| Call Answer Timeliness (cat) | Y | | | 89.92% | R | |
| Utilization | | | | | | |
| Frequency of Ongoing Prenatal Care (fpc) | Y | | | | | |
| <i><21 Percent</i> | | | | 7.74% | R | |
| <i>21-40 Percent</i> | | | | 5.97% | R | |
| <i>41-60 Percent</i> | | | | 9.29% | R | |
| <i>61-80 Percent</i> | | | | 16.15% | R | |
| <i>81+ Percent</i> | | | | 60.84% | R | |
| Well-Child Visits in the First 15 Months of Life (w15) | Y | | Y | | | |
| <i>0 Visits</i> | | | | 1.55% | R | |
| <i>1 Visit</i> | | | | 0.66% | R | |
| <i>2 Visits</i> | | | | 1.77% | R | |
| <i>3 Visits</i> | | | | 3.98% | R | |
| <i>4 Visits</i> | | | | 8.19% | R | |
| <i>5 Visits</i> | | | | 20.80% | R | |
| <i>6+ Visits</i> | | | | 63.05% | R | |
| Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (w34) | Y | | N | 58.28% | R | |
| Adolescent Well-Care Visits (awc) | Y | | N | 35.87% | R | |
| Frequency of Selected Procedures (fsp) | Y | | | | R | |
| Ambulatory Care: Total (amba) | Y | | | | R | |
| Ambulatory Care: Dual Eligibles (ambb) | Y | | | | R | |
| Ambulatory Care: Disabled (ambc) | Y | | | | R | |
| Ambulatory Care: Other (ambd) | Y | | | | R | |
| Inpatient Utilization--General Hospital/Acute Care: Total (ipua) | Y | | | | R | |
| Inpatient Utilization--General Hospital/Acute Care: Dual Eligibles (ipub) | Y | | | | R | |
| Inpatient Utilization--General Hospital/Acute Care: Disabled (ipuc) | Y | | | | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| Inpatient Utilization--General Hospital/Acute Care: Other (ipud) | Y | | | | R | |
| Identification of Alcohol and Other Drug Services: Total (iada) | N | N | | | NR | |
| Identification of Alcohol and Other Drug Services: Dual Eligibles (iadb) | N | N | | | NR | |
| Identification of Alcohol and Other Drug Services: Disabled (iadc) | N | N | | | NR | |
| Identification of Alcohol and Other Drug Services: Other (iadd) | N | N | | | NR | |
| Mental Health Utilization: Total (mpta) | N | N | | | NR | |
| Mental Health Utilization: Dual Eligibles (mptb) | N | N | | | NR | |
| Mental Health Utilization: Disabled (mptc) | N | N | | | NR | |
| Mental Health Utilization: Other (mptd) | N | N | | | NR | |
| Antibiotic Utilization: Total (abxa) | Y | Y | | | R | |
| Antibiotic Utilization: Dual Eligibles (abxb) | Y | Y | | | R | |
| Antibiotic Utilization: Disabled (abxc) | Y | Y | | | R | |
| Antibiotic Utilization: Other (abxd) | Y | Y | | | R | |
| Relative Resource Use | | | | | | |
| Relative Resource Use for People With Diabetes (rdi) | Y | | | | R | |
| Relative Resource Use for People With Asthma (ras) | Y | Y | | | R | |
| Relative Resource Use for People With Cardiovascular Conditions (rca) | Y | | | | R | |
| Relative Resource Use for People With Hypertension (rhy) | Y | | | | R | |
| Relative Resource Use for People With COPD (rco) | Y | | | | R | |
| Health Plan Descriptive Information | | | | | | |
| Board Certification (bcr) | N | | | | NR | |
| Total Membership (tlm) | Y | | | | R | |
| Enrollment by Product Line: Total (enpa) | Y | | | | R | |

| Audit Review Table | | | | | | |
|---|----------------|-----------------|-----------------|------|------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012 | | | | | | |
| This submission is unlocked. | | | | | | |
| Pharmacy Reversal - Unknown | | | | | | |
| Measure/Data Element | Report Measure | Benefit Offered | Rotated Measure | Rate | Reportable | Comment |
| Enrollment by Product Line: Dual Eligibles (enpb) | N | | | | NR | |
| Enrollment by Product Line: Disabled (enpc) | N | | | | NR | |
| Enrollment by Product Line: Other (enpd) | N | | | | NR | |
| Enrollment by State (ebs) | Y | | | | R | |
| Race/Ethnicity Diversity of Membership (rdm) | N | | | | NR | |
| Language Diversity of Membership (ldm) | N | | | | NR | |
| Weeks of Pregnancy at Time of Enrollment in MCO (wop) | Y | | | | R | |

| Adult BMI Assessment (ABA) | |
|--|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | H |
| Eligible population | 1494 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 37 |
| Current year's administrative rate (before exclusions) | 2.48% |
| Minimum required sample size (MRSS) or other sample size | 411 |
| Oversampling rate | 0.1 |
| Final sample size (FSS) | 453 |
| Number of numerator events by administrative data in FSS | 11 |
| Administrative rate on FSS | 2.43% |
| Number of original sample records excluded because of valid data errors | 0 |
| Number of administrative data records excluded | 0 |
| Number of medical records excluded | 0 |
| Number of employee/dependent medical records excluded | 3 |
| Records added from the oversample list | 0 |
| Denominator | 450 |
| Numerator events by administrative data | 11 |
| Numerator events by medical records | 315 |
| Reported rate | 72.44% |
| Lower 95% confidence interval | 68.21% |
| Upper 95% confidence interval | 76.68% |

| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) | | | | | | | | | |
|--|----------------|-------------|--------|--------------------------|-------------|--------|----------------------------------|-------------|--------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | | |
| Data Element | BMI Percentile | | | Counseling for Nutrition | | | Counseling for Physical Activity | | |
| | 3-11 years | 12-17 years | Total | 3-11 years | 12-17 years | Total | 3-11 years | 12-17 years | Total |
| Measurement year | 2012 | | | 2012 | | | 2012 | | |
| Data collection methodology (administrative or hybrid) | H | | | H | | | H | | |
| Eligible population | 5442 | 2366 | 7,808 | 5442 | 2366 | 7,808 | 5442 | 2366 | 7,808 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 97 | 65 | 162 | 116 | 56 | 172 | 48 | 23 | 71 |
| Current year's administrative rate (before exclusions) | 1.78% | 2.75% | 2.07% | 2.13% | 2.37% | 2.20% | 0.88% | 0.97% | 0.91% |
| Minimum required sample size (MRSS) or other sample size | 411 | | | 411 | | | 411 | | |
| Oversampling rate | 0.1 | | | 0.1 | | | 0.1 | | |
| Final sample size | 453 | | | 453 | | | 453 | | |
| Number of numerator events by administrative data in FSS | 6 | 4 | 10 | 6 | 0 | 6 | 2 | 0 | 2 |
| Administrative rate on FSS | 1.32% | 0.88% | 2.21% | 1.32% | 0.00% | 1.32% | 0.44% | 0.00% | 0.44% |
| Number of original sample records excluded because of valid data errors | 0 | | | 0 | | | 0 | | |
| Number of administrative data records excluded | 0 | | | 0 | | | 0 | | |
| Number of medical records excluded | 0 | | | 0 | | | 0 | | |
| Number of employee/dependent medical records excluded | 4 | | | 4 | | | 4 | | |
| Records added from the oversample list | 0 | | | 0 | | | 0 | | |
| Denominator | 304 | 145 | 449 | 304 | 145 | 449 | 304 | 145 | 449 |
| Numerator events by administrative data | 6 | 4 | 10 | 6 | 0 | 6 | 2 | 0 | 2 |
| Numerator events by medical records | 139 | 62 | 201 | 135 | 45 | 180 | 78 | 41 | 119 |
| Reported rate | 47.70% | 45.52% | 46.99% | 46.38% | 31.03% | 41.43% | 26.32% | 28.28% | 26.95% |
| Lower 95% confidence interval | 41.92% | 37.07% | 42.27% | 40.61% | 23.16% | 36.76% | 21.20% | 20.60% | 22.73% |
| Upper 95% confidence interval | 53.48% | 53.97% | 51.72% | 52.15% | 38.91% | 46.09% | 31.43% | 35.95% | 31.16% |

| Childhood Immunization Status (CIS) | | | | | | | | | | | | | | | | | | | | |
|--|----------------------|--------|--------|--------|--------|-------------|--------|-------------------------|-------------|-----------|-----------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | | | | | | | | | | | | | |
| Data Element | General Measure Data | DTaP | IPV | MMR | HiB | Hepatitis B | VZV | Pneumo-coccal Conjugate | Hepatitis A | Rotavirus | Influenza | Combinatio n 2 | Combinatio n 3 | Combinatio n 4 | Combinatio n 5 | Combinatio n 6 | Combinatio n 7 | Combinatio n 8 | Combinatio n 9 | Combinatio n 10 |
| Measurement year | 2012 | | | | | | | | | | | | | | | | | | | |
| Data collection methodology (administrative or hybrid) | H | | | | | | | | | | | | | | | | | | | |
| Eligible population | 1183 | | | | | | | | | | | | | | | | | | | |
| Number of numerator events by admin data in eligible population (before exclusions) | | 436 | 595 | 797 | 684 | 151 | 807 | 440 | 817 | 445 | 434 | 102 | 91 | 84 | 73 | 48 | 70 | 46 | 40 | 39 |
| Current year's administrative rate (before exclusions) | | 36.86% | 50.30% | 67.37% | 57.82% | 12.76% | 68.22% | 37.19% | 69.06% | 37.62% | 36.69% | 8.62% | 7.69% | 7.10% | 6.17% | 4.06% | 5.92% | 3.89% | 3.38% | 3.30% |
| Minimum required sample size (MRSS) or other sample Size | 411 | | | | | | | | | | | | | | | | | | | |
| Oversampling rate | 0.1 | | | | | | | | | | | | | | | | | | | |
| Final sample size | 453 | | | | | | | | | | | | | | | | | | | |
| Number of numerator events by admin data in FSS | | 177 | 243 | 316 | 280 | 75 | 320 | 185 | 322 | 170 | 167 | 52 | 47 | 43 | 36 | 23 | 35 | 21 | 19 | 18 |
| Administrative rate on FSS | | 39.07% | 53.64% | 69.76% | 61.81% | 16.56% | 70.64% | 40.84% | 71.08% | 37.53% | 36.87% | 11.48% | 10.38% | 9.49% | 7.95% | 5.08% | 7.73% | 4.64% | 4.19% | 3.97% |
| Number of original records excluded because of valid data errors | 0 | | | | | | | | | | | | | | | | | | | |
| Number of administrative data records excluded | 0 | | | | | | | | | | | | | | | | | | | |
| Number of medical data records excluded | 0 | | | | | | | | | | | | | | | | | | | |
| Number of employee/dependent medical records excluded | 0 | | | | | | | | | | | | | | | | | | | |
| Records added from the oversample list | 0 | | | | | | | | | | | | | | | | | | | |
| Denominator | 453 | | | | | | | | | | | | | | | | | | | |
| Numerator events by administrative data | | 177 | 243 | 316 | 280 | 75 | 320 | 185 | 322 | 170 | 167 | 52 | 47 | 43 | 36 | 23 | 35 | 21 | 19 | 18 |
| Number of numerator events by medical records | | 200 | 183 | 103 | 145 | 339 | 94 | 196 | 83 | 171 | 76 | 308 | 300 | 295 | 260 | 189 | 254 | 186 | 169 | 166 |
| Reported rate | | 83.22% | 94.04% | 92.49% | 93.82% | 91.39% | 91.39% | 84.11% | 89.40% | 75.28% | 53.64% | 79.47% | 76.60% | 74.61% | 65.34% | 46.80% | 63.80% | 45.70% | 41.50% | 40.62% |
| Lower 95% confidence interval | | 79.67% | 91.75% | 89.96% | 91.49% | 88.70% | 88.70% | 80.63% | 86.46% | 71.19% | 48.94% | 75.64% | 72.59% | 70.50% | 60.85% | 42.09% | 59.26% | 41.00% | 36.85% | 35.99% |
| Upper 95% confidence interval | | 86.77% | 96.33% | 95.03% | 96.15% | 94.08% | 94.08% | 87.58% | 92.35% | 79.36% | 58.34% | 83.30% | 80.61% | 78.73% | 69.83% | 51.50% | 68.33% | 50.39% | 46.15% | 45.25% |

| Immunizations for Adolescents (IMA) | | | | |
|--|----------------------|---------------|---------|---------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | |
| Data Element | General Measure Data | Meningococcal | Tdap/Td | Combination 1 |
| Measurement year | NR | | | |
| Data collection methodology (administrative or hybrid) | NR | | | |
| Eligible population | NR | | | |
| Number of numerator events by admin data in eligible population (before exclusions) | | NR | NR | NR |
| Current year's administrative rate (before exclusions) | | NR | NR | NR |
| Minimum required sample size (MRSS) or other sample Size | NR | | | |
| Oversampling rate | NR | | | |
| Final sample size | NR | | | |
| Number of numerator events by admin data in FSS | | NR | NR | NR |
| Administrative rate on FSS | | NR | NR | NR |
| Number of original records excluded because of valid data errors | NR | | | |
| Number of administrative data records excluded | NR | | | |
| Number of medical data records excluded | NR | | | |
| Number of employee/dependent medical records excluded | NR | | | |
| Records added from the oversample list | NR | | | |
| Denominator | NR | | | |
| Numerator events by administrative data | | NR | NR | NR |
| Number of numerator events by medical records | | NR | NR | NR |
| Reported rate | | NR | NR | NR |
| Lower 95% confidence interval | | NR | NR | NR |
| Upper 95% confidence interval | | NR | NR | NR |

| Human Papillomavirus Vaccine for Female Adolescents (HPV) | |
|--|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | A |
| Eligible population | 240 |
| Number of numerator events by administrative data in eligible population (before exclusions) | NR |
| Current year's administrative rate (before exclusions) | NR |
| Minimum required sample size (MRSS) or other sample size | NR |
| Oversampling rate | NR |
| Final sample size (FSS) | NR |
| Number of numerator events by administrative data in FSS | NR |
| Administrative rate on FSS | NR |
| Number of original sample records excluded because of valid data errors | NR |
| Number of administrative data records excluded | NR |
| Number of medical data records excluded | NR |
| Number of employee/dependent medical records excluded | NR |
| Records added from the oversample list | NR |
| Denominator | NR |
| Numerator events by administrative data | 22 |
| Numerator events by medical records | NR |
| Reported rate | 9.17% |
| Lower 95% confidence interval | 5.31% |
| Upper 95% confidence interval | 13.03% |

| Lead Screening in Children (LSC) | |
|--|-----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Lead Screening in Children | |
| Data Elements which do not apply to the selected data collection methodology will not appear | General Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | A |
| Eligible population | 1183 |
| Number of numerator events by admin data in eligible population (before exclusions) | NR |
| Current year's administrative rate (before exclusions) | NR |
| Minimum required sample size (MRSS) or other sample size | NR |
| Oversampling rate | NR |
| Final sample size (FSS) | NR |
| Number of numerator events by administrative data in FSS | NR |
| Administrative rate on FSS | NR |
| Number of original sample records excluded because of valid data errors | NR |
| Number of administrative data records excluded | NR |
| Number of medical data records excluded | NR |
| Number of employee/dependent medical records excluded | NR |
| Records added from the oversample list | NR |
| Denominator | NR |
| Numerator events by administrative data | 403 |
| Numerator events by medical records | NR |
| Reported rate | 34.07% |
| Lower 95% confidence interval | 31.32% |
| Upper 95% confidence interval | 36.81% |

| Breast Cancer Screening (BCS) | |
|--|-----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | General Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 277 |
| Numerator events by administrative data | 154 |
| Reported rate | 55.60% |
| Lower 95% confidence interval | 49.56% |
| Upper 95% confidence interval | 61.63% |

| Cervical Cancer Screening (CCS) | |
|--|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | H |
| Eligible population | 1857 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 974 |
| Current year's administrative rate (before exclusions) | 52.45% |
| Minimum required sample size (MRSS) or other sample size | 411 |
| Oversampling rate | 0.1 |
| Final sample size (FSS) | 453 |
| Number of numerator events by administrative data in FSS | 231 |
| Administrative rate on FSS | 50.99% |
| Number of original sample records excluded because of valid data errors | 1 |
| Number of administrative data records excluded | 0 |
| Number of medical data records excluded | 2 |
| Number of employee/dependent medical records excluded | 0 |
| Records added from the oversample list | 0 |
| Denominator | 450 |
| Numerator events by administrative data | 231 |
| Numerator events by medical records | 32 |
| Reported rate | 58.44% |
| Lower 95% confidence interval | 53.78% |
| Upper 95% confidence interval | 63.11% |

| Chlamydia Screening in Women (CHL) | | | | |
|--|----------------------|-------------|-------------|--------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | |
| Data Element | General Measure Data | 16-20 years | 21-24 years | Total |
| Measurement year | 2012 | | | |
| Data collection methodology (administrative) | A | | | |
| Eligible population | | 458 | 292 | 750 |
| Numerator events by administrative data | | 185 | 159 | 344 |
| Reported rate | | 40.39% | 54.45% | 45.87% |
| Lower 95% confidence interval | | 35.79% | 48.57% | 42.23% |
| Upper 95% confidence interval | | 45.00% | 60.34% | 49.50% |

Appropriate Testing for Children with Pharyngitis (CWP)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|---------------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 775 |
| Numerator events by administrative data | 492 |
| Reported rate | 63.48% |
| Lower 95% confidence interval | 60.03% |
| Upper 95% confidence interval | 66.94% |

| Appropriate Treatment for Children With URI (URI) | |
|--|--|
|--|--|

| | |
|---|--|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
|---|--|

| Data Element | Measure Data |
|---|---------------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 1868 |
| Numerator events by administrative data | 281 |
| Reported rate | 84.96% |
| Lower 95% confidence interval | 83.31% |
| Upper 95% confidence interval | 86.61% |

| Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB) | |
|--|---------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 91 |
| Total numerator events by administrative data | 75 |
| Reported rate | 17.58% |
| Lower 95% confidence interval | 9.21% |
| Upper 95% confidence interval | 25.95% |

Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|--------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 15 |
| Numerator events by administrative data | 2 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

| Pharmacotherapy Management of COPD Exacerbation (PCE) | | | |
|--|---------------------|--------------------------------|-----------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Pharmacotherapy Management of COPD Exacerbation | | | |
| Data Elements | Measure Data | Systemic corticosteroid | Bronchodilator |
| Measurement year | 2012 | | |
| Data collection methodology (administrative) | A | | |
| Eligible population | 25 | | |
| Numerator events by administrative data | | 19 | 23 |
| Reported rate | | NA | NA |
| Lower 95% confidence interval | | NA | NA |
| Upper 95% confidence interval | | NA | NA |

| Use of Appropriate Medications for People With Asthma (ASM) | | | | | | |
|--|-----------------------------|-------------------|--------------------|--------------------|--------------------|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | |
| Data Element | General Measure Data | 5-11 years | 12-18 years | 19-50 years | 51-64 years | Total |
| Measurement year | 2012 | | | | | |
| Data collection methodology (administrative) | A | | | | | |
| Eligible population | | 82 | 63 | 43 | 4 | 192 |
| Number of required exclusions | | 5 | 4 | 13 | 22 | 44 |
| Numerator events by administrative data | | 77 | 49 | 37 | 3 | 166 |
| Reported rate | | 93.90% | 77.78% | 86.05% | NA | 86.46% |
| Lower 95% confidence interval | | 88.11% | 66.72% | 74.53% | NA | 81.36% |
| Upper 95% confidence interval | | 99.69% | 88.84% | 97.57% | NA | 91.56% |

| Medication Management for People With Asthma (MMA) | | | | | | | | | | | |
|--|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | | | | |
| Data Element | General Measure Data | 5-11 years | | 12-18 years | | 19-50 years | | 51-64 years | | Total | |
| | | Medication Compliance 50% | Medication Compliance 75% | Medication Compliance 50% | Medication Compliance 75% | Medication Compliance 50% | Medication Compliance 75% | Medication Compliance 50% | Medication Compliance 75% | Medication Compliance 50% | Medication Compliance 75% |
| Measurement year | 2012 | | | | | | | | | | |
| Data collection methodology (administrative) | A | | | | | | | | | | |
| Eligible population | | 77 | | 49 | | 37 | | 3 | | 166 | |
| Number of required exclusions | | 10 | | 18 | | 19 | | 23 | | 70 | |
| Numerator events by administrative data | | 34 | 18 | 16 | 8 | 17 | 11 | 3 | 1 | 70 | 38 |
| Reported rate | | 44.16% | 23.38% | 32.65% | 16.33% | 45.95% | 29.73% | NA | NA | 42.17% | 22.89% |
| Lower 95% confidence interval | | 32.41% | 13.27% | 18.50% | 4.96% | 28.54% | 13.65% | NA | NA | 34.36% | 16.20% |
| Upper 95% confidence interval | | 55.90% | 33.48% | 46.80% | 27.70% | 63.36% | 45.81% | NA | NA | 49.98% | 29.58% |

| Asthma Medication Ratio (AMR) | | | | | | |
|--|--------------|------------|-------------|-------------|-------------|--------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | |
| Data Element | Measure Data | 5-11 years | 12-18 years | 19-50 years | 51-64 years | Total |
| Measurement year | 2012 | | | | | |
| Data collection methodology (administrative) | A | | | | | |
| Eligible population | | 82 | 63 | 43 | 4 | 192 |
| Number of required exclusions | | 5 | 4 | 13 | 22 | 44 |
| Numerator events by administrative data | | 54 | 26 | 21 | 3 | 104 |
| Reported rate | | 65.85% | 41.27% | 48.84% | NA | 54.17% |
| Lower 95% confidence interval | | 54.98% | 28.32% | 32.73% | NA | 46.86% |
| Upper 95% confidence interval | | 76.73% | 54.22% | 64.94% | NA | 61.48% |

Cholesterol Management for Patients With Cardiovascular Conditions (CMC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | General Measure Data | LDL-C Screening | LDL-C level <100 mg/dL |
|--|----------------------|-----------------|------------------------|
| Measurement year | 2012 | | |
| Data collection methodology (administrative or hybrid) | H | | |
| Eligible population | 17 | | |
| Number of numerator events by administrative data in eligible population (before exclusions) | | 10 | 0 |
| Current year's administrative rate (before exclusions) | | 58.82% | 0.00% |
| Minimum required sample size (MRSS) or other sample size | 17 | | |
| Oversampling rate | 0.1 | | |
| Final sample size (FSS) | 17 | | |
| Number of numerator events by administrative data in FSS | | 10 | 0 |
| Administrative rate on FSS | | 58.82% | 0.00% |
| Number of original sample records excluded because of valid data errors | 0 | | |
| Number of employee/dependent medical records excluded | 0 | | |
| Records added from the oversample list | 0 | | |
| Denominator | 17 | | |
| Numerator events by administrative data | | 10 | 0 |
| Numerator events by medical records | | 0 | 3 |
| Reported rate | | NA | NA |
| Lower 95% confidence interval | | NA | NA |
| Upper 95% confidence interval | | NA | NA |

| Controlling High Blood Pressure (CBP) | |
|--|-----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | General Measure Data |
| Measurement year | 2012 |
| Data collection methodology (hybrid) | H |
| Eligible population | 344 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 0 |
| Current year's administrative rate (before exclusions) | 0.00% |
| Minimum required sample size (MRSS) or other sample size | 326 |
| Oversampling rate | 0.15 |
| Final sample size (FSS) | 326 |
| Number of numerator events by administrative data in FSS | 0 |
| Administrative rate on FSS | 0.00% |
| Number of original sample records excluded because of valid data errors | 6 |
| Number of records excluded because of false positive diagnoses | 6 |
| Number of administrative data records excluded | 2 |
| Number of medical data records excluded | 0 |
| Number of employee/dependent medical records excluded | 0 |
| Records added from the oversample list | 0 |
| Denominator | 318 |
| Numerator events by administrative data | 0 |
| Numerator events by medical records | 156 |
| Reported rate | 49.06% |
| Lower 95% confidence interval | 43.40% |
| Upper 95% confidence interval | 54.71% |

| Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) | |
|--|---------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 0 |
| Numerator events by administrative data | 0 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

| Comprehensive Diabetes Care (CDC) | | | | | | | | | | |
|--|---------------|----------------------------|-----------------------|---|----------|-----------------|------------------------|-----------------------------------|---|---|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | | | |
| Data Element | HbA1c Testing | HbA1c Poor Control (>9.0%) | HbA1c Control (<8.0%) | HbA1c Control (<7.0%) for a Selected Population | Eye Exam | LDL-C Screening | LDL-C Level <100 mg/dL | Medical Attention for Nephropathy | Blood Pressure Controlled <140/80 mm Hg | Blood Pressure Controlled <140/90 mm Hg |
| Measurement year | 2011 | 2011 | 2011 | 2011 | 2011 | 2011 | 2011 | 2011 | 2011 | 2011 |
| Data collection methodology (administrative or hybrid) | H | H | H | NR | H | H | H | H | H | H |
| Eligible population | 240 | 240 | 240 | NR | 240 | 240 | 240 | 240 | 240 | 240 |
| Number of numerator events by administrative data in eligible population (before optional exclusions) | 177 | 228 | 11 | NR | 105 | 159 | 18 | 167 | 0 | 0 |
| Current year's administrative rate (before optional exclusions) | 73.75% | 95.00% | 4.58% | NR | 43.75% | 66.25% | 7.50% | 69.58% | 0.00% | 0.00% |
| Minimum required sample size (MRSS) or other sample size | 235 | 235 | 235 | NR | 235 | 235 | 235 | 235 | 235 | 235 |
| Oversampling rate | 0.1 | 0.1 | 0.1 | NR | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Final sample size (FSS) | 235 | 235 | 235 | NR | 235 | 235 | 235 | 235 | 235 | 235 |
| Number of numerator events by administrative data in FSS | 176 | 223 | 11 | NR | 105 | 159 | 16 | 163 | 0 | 0 |
| Administrative rate on FSS | 74.89% | 94.89% | 4.68% | NR | 44.68% | 67.66% | 6.81% | 69.36% | 0.00% | 0.00% |
| Number of original sample records excluded because of valid data errors | 3 | 3 | 3 | NR | 3 | 3 | 3 | 3 | 3 | 3 |
| Number of optional administrative data records excluded | 0 | 0 | 0 | NR | 0 | 0 | 0 | 0 | 0 | 0 |
| Number of optional medical data records excluded | 0 | 0 | 0 | NR | 0 | 0 | 0 | 0 | 0 | 0 |
| Number of HbA1c <7 required medical records excluded | | | | NR | | | | | | |
| Number of HbA1c <7 required administrative data records excluded | | | | NR | | | | | | |
| Number of employee/dependent medical records excluded | 0 | 0 | 0 | NR | 0 | 0 | 0 | 0 | 0 | 0 |
| Records added from the oversample list | 0 | 0 | 0 | NR | 0 | 0 | 0 | 0 | 0 | 0 |
| Denominator | 232 | 232 | 232 | NR | 232 | 232 | 232 | 232 | 232 | 232 |
| Numerator events by administrative data | 176 | 59 | 11 | NR | 105 | 159 | 16 | 163 | 0 | 0 |
| Numerator events by medical records | 15 | 33 | 95 | NR | 20 | 9 | 75 | 14 | 108 | 151 |
| Reported rate | 82.33% | 39.66% | 45.69% | NR | 53.88% | 72.41% | 39.22% | 76.29% | 46.55% | 65.09% |
| Lower 95% confidence interval | 77.20% | 33.14% | 39.06% | NR | 47.25% | 66.45% | 32.73% | 70.61% | 39.92% | 58.74% |
| Upper 95% confidence interval | 87.45% | 46.17% | 52.32% | NR | 60.51% | 78.38% | 45.72% | 81.98% | 53.19% | 71.44% |

| Disease Modifying Anti-Rheumatic Drug therapy in Rheumatoid Arthritis (ART) | |
|---|---------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 18 |
| Numerator events by administrative data | 14 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

Use of Imaging Studies for Low Back Pain (LBP)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|---------------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 144 |
| Numerator events by administrative data | 29 |
| Reported rate | 79.86% |
| Lower 95% confidence Interval | 72.96% |
| Upper 95% confidence Interval | 86.76% |

| Antidepressant Medication Management (AMM) | | | |
|--|-----------------------------|--|---|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Data Element | General Measure Data | Effective Acute Phase Treatment | Effective Continuation Phase Treatment |
| Measurement year | NR | | |
| Data collection methodology (administrative) | NR | | |
| Eligible population | NR | | |
| Numerator events by administrative data | | NR | NR |
| Reported rate | | NR | NR |
| Lower 95% confidence interval | | NR | NR |
| Upper 95% confidence interval | | NR | NR |

| Follow-Up Care for Children Prescribed ADHD Medication (ADD) | | | |
|--|-----------------------------|-------------------------|---|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Data Element | General Measure Data | Initiation Phase | Continuation and Maintenance Phase |
| Measurement year | 2012 | | |
| Data collection methodology (administrative) | A | | |
| Eligible population | | 76 | 9 |
| Numerator events by administrative data | | 34 | 4 |
| Reported rate | | 44.74% | NA |
| Lower 95% confidence interval | | 32.90% | NA |
| Upper 95% confidence interval | | 56.57% | NA |

| Follow-Up After Hospitalization for Mental Illness (FUH) | | | |
|--|-----------------------------|-------------------------|------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Data Element | General Measure Data | 30-day follow-up | 7-day follow-up |
| Measurement year | NR | | |
| Data collection methodology (administrative) | NR | | |
| Eligible population | NR | | |
| Numerator events by administrative data | | NR | NR |
| Reported rate | | NR | NR |
| Lower 95% confidence interval | | NR | NR |
| Upper 95% confidence interval | | NR | NR |

Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (SSD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|--------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 18 |
| Number of required exclusions | 34 |
| Numerator events by administrative data | 17 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

Diabetes Monitoring for People With Diabetes and Schizophrenia (SMD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|---------------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 4 |
| Number of optional exclusions | 0 |
| Numerator events by administrative data | 1 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (SMC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|--------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 0 |
| Numerator events by administrative data | 0 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

**Adherence to Antipsychotic Medications
for Individuals With Schizophrenia
(SAA)**

Blue Cross and Blue Shield of New Mexico (Org ID:
1825, SubID: 9522, Medicaid, Spec Area: None, Spec
Proj: None)

| Data Element | Measure Data |
|---|--------------|
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 10 |
| Number of required exclusions | 17 |
| Numerator events by administrative data | 3 |
| Reported rate | NA |
| Lower 95% confidence interval | NA |
| Upper 95% confidence interval | NA |

| Annual Monitoring for Patients on Persistent Medications (MPM) | | | | | | |
|--|----------------------|------------------------|---------|-----------|------------------|--------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | |
| Data Element | General Measure Data | ACE Inhibitors or ARBs | Digoxin | Diuretics | Anti-convulsants | Total |
| Measurement year | 2012 | | | | | |
| Data collection methodology (administrative) | A | | | | | |
| Eligible population | | 234 | 4 | 146 | 34 | 418 |
| Numerator events by administrative data | | 209 | 4 | 126 | 21 | 360 |
| Reported rate | | 89.32% | NA | 86.30% | 61.76% | 86.12% |
| Lower 95% confidence interval | | 85.14% | NA | 80.38% | 43.96% | 82.69% |
| Upper 95% confidence interval | | 93.49% | NA | 92.22% | 79.57% | 89.56% |

| Adults' Access to Preventive/Ambulatory Health Services (AAP) | | | | | |
|--|----------------------|-------------|-------------|-----------|--------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | |
| Data Element | General Measure Data | 20-44 years | 45-64 years | 65+ years | Total |
| Measurement year | 2012 | | | | |
| Data collection methodology (administrative) | A | | | | |
| Eligible population | | 2487 | 869 | 1 | 3,357 |
| Numerator events by administrative data | | 2032 | 739 | 1 | 2,772 |
| Reported rate | | 81.70% | 85.04% | NA | 82.57% |
| Lower 95% confidence interval | | 80.17% | 82.61% | NA | 81.28% |
| Upper 95% confidence interval | | 83.24% | 87.47% | NA | 83.87% |

| Children and Adolescents' Access to Primary Care Practitioners (CAP) | | | | | |
|--|----------------------|--------------|---------------------|------------|-------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | |
| Data Element | General Measure Data | 12-24 months | 25 months - 6 years | 7-11 years | 12-19 years |
| Measurement year | 2012 | | | | |
| Data collection methodology (administrative) | A | | | | |
| Eligible population | | 1215 | 4964 | 2159 | 2550 |
| Numerator events by administrative data | | 1181 | 4275 | 1840 | 2187 |
| Reported rate | | 97.20% | 86.12% | 85.22% | 85.76% |
| Lower 95% confidence interval | | 96.23% | 85.15% | 83.70% | 84.39% |
| Upper 95% confidence interval | | 98.17% | 87.09% | 86.74% | 87.14% |

| Annual Dental Visit (ADV) | | | | | | | | |
|---|---------------------|------------------|------------------|-------------------|--------------------|--------------------|--------------------|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | |
| Data Element | Measure Data | 2-3 Years | 4-6 Years | 7-10 Years | 11-14 Years | 15-18 Years | 19-21 Years | Total |
| Measurement year | 2012 | | | | | | | |
| Data collection methodology (administrative) | A | | | | | | | |
| Eligible population | | 2490 | 2577 | 2781 | 2466 | 2068 | 259 | 12,641 |
| Numerator events by administrative data | | 1388 | 1766 | 2040 | 1632 | 1167 | 97 | 8,090 |
| Reported rate | | 55.74% | 68.53% | 73.35% | 66.18% | 56.43% | 37.45% | 64.00% |
| Lower 95% confidence interval | | 53.77% | 66.72% | 71.69% | 64.29% | 54.27% | 31.36% | 63.16% |
| Upper 95% confidence interval | | 57.71% | 70.34% | 75.02% | 68.07% | 58.59% | 43.54% | 64.84% |

| Initiation and Engagement of AOD Dependence Treatment (IET) | | | | | | | |
|--|----------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | |
| Data Elements | General Measure Data | 13-17 years | | 18+ years | | Total | |
| | | Initiation of AOD Treatment | Engagement of AOD Treatment | Initiation of AOD Treatment | Engagement of AOD Treatment | Initiation of AOD Treatment | Engagement of AOD Treatment |
| Measurement year | NR | | | | | | |
| Data collection methodology (administrative) | NR | | | | | | |
| Eligible population | | NR | | NR | | NR | |
| Numerator events by administrative data | | NR | NR | NR | NR | NR | NR |
| Reported rate | | NR | NR | NR | NR | NR | NR |
| Lower 95% confidence interval | | NR | NR | NR | NR | NR | NR |
| Upper 95% confidence interval | | NR | NR | NR | NR | NR | NR |

| Prenatal and Postpartum Care (PPC) | | |
|---|------------------------------------|------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | |
| Data Element | Timeliness of Prenatal Care | Postpartum Care |
| Measurement year | 2012 | 2012 |
| Data collection methodology (administrative or hybrid) | H | H |
| Eligible population | 1161 | 1161 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 425 | 379 |
| Current year's administrative rate (before exclusions) | 36.61% | 32.64% |
| Minimum required sample size (MRSS) or other sample size | 411 | 411 |
| Oversampling rate | 0.1 | 0.1 |
| Final sample size (FSS) | 453 | 453 |
| Number of numerator events by administrative data in FSS | 168 | 146 |
| Administrative rate on FSS | 37.09% | 32.23% |
| Number of original sample records excluded because of valid data errors | 1 | 1 |
| Number of employee/dependent medical records excluded | 0 | 0 |
| Records added from the oversample list | 0 | 0 |
| Denominator | 452 | 452 |
| Numerator events by administrative data | 168 | 146 |
| Numerator events by medical records | 221 | 139 |
| Reported rate | 86.06% | 63.05% |
| Lower 95% confidence interval | 82.76% | 58.49% |
| Upper 95% confidence interval | 89.37% | 67.61% |

| Call Answer Timeliness (CAT) | |
|---|---------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative) | A |
| Eligible population | 19472 |
| Numerator events by administrative data | 17510 |
| Reported rate | 89.92% |
| Lower 95% confidence interval | 89.50% |
| Upper 95% confidence interval | 90.35% |

Frequency of Ongoing Prenatal Care (FPC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | General Measure Data | <21 Percent | 21-40 Percent | 41-60 Percent | 61-80 Percent | 81+ Percent |
|--|----------------------|-------------|---------------|---------------|---------------|-------------|
| Measurement year | 2012 | | | | | |
| Data collection methodology (administrative or hybrid) | H | | | | | |
| Eligible population | 1160 | | | | | |
| Number of numerator events by administrative data in eligible population (before exclusions) | | 737 | 209 | 86 | 54 | 74 |
| Current year's administrative rate (before exclusions) | | 63.53% | 18.02% | 7.41% | 4.66% | 6.38% |
| Minimum required sample size (MRSS) or other sample size | 411 | | | | | |
| Oversampling rate | 0.1 | | | | | |
| Final sample size (FSS) | 453 | | | | | |
| Number of numerator events by administrative data in FSS | | 288 | 89 | 29 | 15 | 32 |
| Administrative rate on FSS | | 63.58% | 19.65% | 6.40% | 3.31% | 7.06% |
| Number of original sample records excluded because of valid data errors | 1 | | | | | |
| Number of employee/dependent medical records excluded | 0 | | | | | |
| Records added from the oversample list | 0 | | | | | |
| Denominator | 452 | | | | | |
| Numerator events by administrative data | | 32 | 18 | 5 | 8 | 36 |
| Numerator events by medical records | | 3 | 9 | 37 | 65 | 239 |
| Reported rate | | 7.74% | 5.97% | 9.29% | 16.15% | 60.84% |
| Lower 95% confidence interval | | 5.17% | 3.68% | 6.50% | 12.65% | 56.23% |
| Upper 95% confidence interval | | 10.32% | 8.27% | 12.08% | 19.65% | 65.45% |

| Well-Child Visits in the First 15 Months of Life (W15) | | | | | | | | |
|--|--------------|----------|---------|----------|----------|----------|----------|------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | | | | | |
| Data Element | Measure Data | 0 visits | 1 visit | 2 visits | 3 visits | 4 visits | 5 visits | 6 or more visits |
| Measurement year | 2011 | | | | | | | |
| Data collection methodology (administrative or hybrid) | H | | | | | | | |
| Eligible population | 938 | | | | | | | |
| Number of numerator events by administrative data in eligible population (before exclusions) | | 18 | 19 | 29 | 75 | 141 | 215 | 441 |
| Current year's administrative rate (before exclusions) | | 1.92% | 2.03% | 3.09% | 8.00% | 15.03% | 22.92% | 47.01% |
| Minimum required sample size (MRSS) or other sample size | 411 | | | | | | | |
| Oversampling rate | 0.1 | | | | | | | |
| Final sample size (FSS) | 453 | | | | | | | |
| Number of numerator events by administrative data in FSS | | 9 | 5 | 11 | 32 | 71 | 110 | 215 |
| Administrative rate on FSS | | 1.99% | 1.10% | 2.43% | 7.06% | 15.67% | 24.28% | 47.46% |
| Number of original sample records excluded because of valid data errors | 0 | | | | | | | |
| Number of employee/dependent medical records excluded | 1 | | | | | | | |
| Records added from the oversample list | 0 | | | | | | | |
| Denominator | 452 | | | | | | | |
| Numerator events by administrative data | | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Numerator events by medical records | | 7 | 3 | 8 | 18 | 37 | 94 | 285 |
| Reported rate | | 1.55% | 0.66% | 1.77% | 3.98% | 8.19% | 20.80% | 63.05% |
| Lower 95% confidence interval | | 0.30% | 0.00% | 0.44% | 2.07% | 5.55% | 16.94% | 58.49% |
| Upper 95% confidence interval | | 2.80% | 1.52% | 3.10% | 5.90% | 10.82% | 24.65% | 67.61% |

| Adolescent Well-Care Visits (AWC) | |
|--|--------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Data Element | Measure Data |
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | A |
| Eligible population | 4179 |
| Number of numerator events by administrative data in eligible population (before exclusions) | NR |
| Current year's administrative rate (before exclusions) | NR |
| Minimum required sample size (MRSS) or other sample size | NR |
| Oversampling rate | NR |
| Final sample size (FSS) | NR |
| Number of numerator events by administrative data in FSS | NR |
| Administrative rate on FSS | NR |
| Number of original sample records excluded because of valid data errors | NR |
| Number of employee/dependent medical records excluded | NR |
| Records added from the oversample list | NR |
| Denominator | NR |
| Numerator events by administrative data | 1499 |
| Numerator events by medical records | NR |
| Reported rate | 35.87% |
| Lower 95% confidence interval | 34.40% |
| Upper 95% confidence interval | 37.34% |

Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (W34)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Data Element | Measure Data |
|--|--------------|
| Measurement year | 2012 |
| Data collection methodology (administrative or hybrid) | H |
| Eligible population | 3778 |
| Number of numerator events by administrative data in eligible population (before exclusions) | 2176 |
| Current year's administrative rate (before exclusions) | 57.60% |
| Minimum required sample size (MRSS) or other sample size | 411 |
| Oversampling rate | 0.1 |
| Final sample size (FSS) | 453 |
| Number of numerator events by administrative data in FSS | 261 |
| Administrative rate on FSS | 57.62% |
| Number of original sample records excluded because of valid data errors | 0 |
| Number of employee/dependent medical records excluded | 0 |
| Records added from the oversample list | 0 |
| Denominator | 453 |
| Numerator events by administrative data | 261 |
| Numerator events by medical records | 3 |
| Reported rate | 58.28% |
| Lower 95% confidence interval | 53.63% |
| Upper 95% confidence interval | 62.93% |

Frequency of Selected Procedures (FSP)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Male | Female | Total | |
|-------|--------|--------|---------|--|
| 0-9 | 82,181 | 77,791 | 159,972 | |
| 10-19 | 44,970 | 45,216 | 90,186 | |
| 15-44 | | 63,210 | | |
| 20-44 | 15,621 | 43,592 | | |
| 30-64 | 17,011 | | | |
| 45-64 | 8,187 | 7,233 | | |

| Procedure | Age | Sex | Number of Procedures | Procedures / 1,000 Member Years |
|--------------------------------|-------|---------------|----------------------|---------------------------------|
| Bariatric weight loss surgery | 0-19 | Male | 0 | 0.00 |
| | | Female | 0 | 0.00 |
| | 20-44 | Male | 0 | 0.00 |
| | | Female | 0 | 0.00 |
| | 45-64 | Male | 0 | 0.00 |
| | | Female | 0 | 0.00 |
| Tonsillectomy | 0-9 | Male & Female | 92 | 0.58 |
| | 10-19 | | 38 | 0.42 |
| Hysterectomy, Abdominal | 15-44 | Female | 10 | 0.16 |
| | 45-64 | | 6 | 0.83 |
| Hysterectomy, Vaginal | 15-44 | Female | 11 | 0.17 |
| | 45-64 | | 2 | 0.28 |
| Cholecystectomy, Open | 30-64 | Male | 2 | 0.12 |
| | 15-44 | Female | 0 | 0.00 |
| | | | 0 | 0.00 |
| Cholecystectomy (laparoscopic) | 30-64 | Male | 6 | 0.35 |
| | 15-44 | Female | 63 | 1.00 |
| | | | 4 | 0.55 |
| Back Surgery | 20-44 | Male | 4 | 0.26 |
| | | Female | 6 | 0.14 |
| | 45-64 | Male | 8 | 0.98 |
| | | Female | 9 | 1.24 |
| Mastectomy | 15-44 | Female | 0 | 0.00 |
| | 45-64 | | 0 | 0.00 |
| Lumpectomy | 15-44 | Female | 7 | 0.11 |
| | 45-64 | | 5 | 0.69 |

| Ambulatory Care: Total (AMBA) | | | | |
|--|-------------------|-----------------------------|---------------|-----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | | |
| Age | Member Months | | | |
| <1 | 23,881 | | | |
| 1-9 | 136,091 | | | |
| 10-19 | 90,186 | | | |
| 20-44 | 59,213 | | | |
| 45-64 | 15,420 | | | |
| 65-74 | 8 | | | |
| 75-84 | 0 | | | |
| 85+ | 0 | | | |
| Unknown | 0 | | | |
| Total | 324,799 | | | |
| Age | Outpatient Visits | | ED Visits | |
| | Visits | Visits/ 1,000 Member Months | Visits | Visits/ 1,000 Member Months |
| <1 | 19146 | 801.73 | 1897 | 79.44 |
| 1-9 | 38633 | 283.88 | 5344 | 39.27 |
| 10-19 | 18964 | 210.28 | 2942 | 32.62 |
| 20-44 | 21897 | 369.80 | 4561 | 77.03 |
| 45-64 | 9647 | 625.62 | 1056 | 68.48 |
| 65-74 | 11 | 1,375.00 | 0 | 0.00 |
| 75-84 | 0 | NA | 0 | NA |
| 85+ | 0 | NA | 0 | NA |
| Unknown | 0 | | 0 | |
| Total | 108,298 | 333.43 | 15,800 | 48.65 |

Ambulatory Care: Dual Eligibles (AMBB)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months | | | | |
|--------------|-------------------|-----------------------------|---------------|-----------------------------|--|
| <1 | 23,881 | | | | |
| 1-9 | 136,091 | | | | |
| 10-19 | 90,186 | | | | |
| 20-44 | 59,213 | | | | |
| 45-64 | 15,420 | | | | |
| 65-74 | 8 | | | | |
| 75-84 | 0 | | | | |
| 85+ | 0 | | | | |
| Unknown | 0 | | | | |
| Total | 324,799 | | | | |
| Age | Outpatient Visits | | ED Visits | | |
| | Visits | Visits/ 1,000 Member Months | Visits | Visits/ 1,000 Member Months | |
| <1 | 19146 | 801.73 | 1897 | 79.44 | |
| 1-9 | 38633 | 283.88 | 5344 | 39.27 | |
| 10-19 | 18964 | 210.28 | 2942 | 32.62 | |
| 20-44 | 21897 | 369.80 | 4561 | 77.03 | |
| 45-64 | 9647 | 625.62 | 1056 | 68.48 | |
| 65-74 | 11 | 1,375.00 | 0 | 0.00 | |
| 75-84 | 0 | NA | 0 | NA | |
| 85+ | 0 | NA | 0 | NA | |
| Unknown | 0 | | 0 | | |
| Total | 108,298 | 333.43 | 15,800 | 48.65 | |

Ambulatory Care: Disabled (AMBC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months | | | | |
|--------------|-------------------|-----------------------------|---------------|-----------------------------|--|
| <1 | 23,881 | | | | |
| 1-9 | 136,091 | | | | |
| 10-19 | 90,186 | | | | |
| 20-44 | 59,213 | | | | |
| 45-64 | 15,420 | | | | |
| 65-74 | 8 | | | | |
| 75-84 | 0 | | | | |
| 85+ | 0 | | | | |
| Unknown | 0 | | | | |
| Total | 324,799 | | | | |
| Age | Outpatient Visits | | ED Visits | | |
| | Visits | Visits/ 1,000 Member Months | Visits | Visits/ 1,000 Member Months | |
| <1 | 19146 | 801.73 | 1897 | 79.44 | |
| 1-9 | 38633 | 283.88 | 5344 | 39.27 | |
| 10-19 | 18964 | 210.28 | 2942 | 32.62 | |
| 20-44 | 21897 | 369.80 | 4561 | 77.03 | |
| 45-64 | 9647 | 625.62 | 1056 | 68.48 | |
| 65-74 | 11 | 1,375.00 | 0 | 0.00 | |
| 75-84 | 0 | NA | 0 | NA | |
| 85+ | 0 | NA | 0 | NA | |
| Unknown | 0 | | 0 | | |
| Total | 108,298 | 333.43 | 15,800 | 48.65 | |

Ambulatory Care: Other (AMBD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months | | | | |
|--------------|-------------------|-----------------------------|---------------|-----------------------------|--|
| <1 | 23,881 | | | | |
| 1-9 | 136,091 | | | | |
| 10-19 | 90,186 | | | | |
| 20-44 | 59,213 | | | | |
| 45-64 | 15,420 | | | | |
| 65-74 | 8 | | | | |
| 75-84 | 0 | | | | |
| 85+ | 0 | | | | |
| Unknown | 0 | | | | |
| Total | 324,799 | | | | |
| Age | Outpatient Visits | | ED Visits | | |
| | Visits | Visits/ 1,000 Member Months | Visits | Visits/ 1,000 Member Months | |
| <1 | 19146 | 801.73 | 1897 | 79.44 | |
| 1-9 | 38633 | 283.88 | 5344 | 39.27 | |
| 10-19 | 18964 | 210.28 | 2942 | 32.62 | |
| 20-44 | 21897 | 369.80 | 4561 | 77.03 | |
| 45-64 | 9647 | 625.62 | 1056 | 68.48 | |
| 65-74 | 11 | 1,375.00 | 0 | 0.00 | |
| 75-84 | 0 | NA | 0 | NA | |
| 85+ | 0 | NA | 0 | NA | |
| Unknown | 0 | | 0 | | |
| Total | 108,298 | 333.43 | 15,800 | 48.65 | |

Inpatient Utilization--General Hospital/Acute Care: Total (IPUA)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months |
|--------------|----------------|
| <1 | 23,881 |
| 1-9 | 136,091 |
| 10-19 | 90,186 |
| 20-44 | 59,213 |
| 45-64 | 15,420 |
| 65-74 | 8 |
| 75-84 | 0 |
| 85+ | 0 |
| Unknown | 0 |
| Total | 324,799 |

Total Inpatient

| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| <1 | 222 | 9.30 | 1328 | 55.61 | 5.98 |
| 1-9 | 313 | 2.30 | 1044 | 7.67 | 3.34 |
| 10-19 | 384 | 4.26 | 1221 | 13.54 | 3.18 |
| 20-44 | 1485 | 25.08 | 4188 | 70.73 | 2.82 |
| 45-64 | 294 | 19.07 | 1727 | 112.00 | 5.87 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 2,698 | 8.31 | 9,508 | 29.27 | 3.52 |

Medicine

| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| <1 | 189 | 7.91 | 753 | 31.53 | 3.98 |
| 1-9 | 261 | 1.92 | 817 | 6.00 | 3.13 |
| 10-19 | 118 | 1.31 | 533 | 5.91 | 4.52 |
| 20-44 | 227 | 3.83 | 1022 | 17.26 | 4.50 |
| 45-64 | 214 | 13.88 | 1084 | 70.30 | 5.07 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 1,009 | 3.11 | 4,209 | 12.96 | 4.17 |

Surgery

| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
|--------------|------------|----------------------------------|--------------|-----------------------------|------------------------|
| <1 | 33 | 1.38 | 575 | 24.08 | 17.42 |
| 1-9 | 52 | 0.38 | 227 | 1.67 | 4.37 |
| 10-19 | 39 | 0.43 | 175 | 1.94 | 4.49 |
| 20-44 | 82 | 1.38 | 349 | 5.89 | 4.26 |
| 45-64 | 77 | 4.99 | 633 | 41.05 | 8.22 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 283 | 0.87 | 1,959 | 6.03 | 6.92 |

Maternity*

| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| 10-19 | 227 | 2.52 | 513 | 5.69 | 2.26 |
| 20-44 | 1176 | 19.86 | 2817 | 47.57 | 2.40 |
| 45-64 | 3 | 0.19 | 10 | 0.65 | 3.33 |
| Unknown | 0 | | 0 | | NA |
| Total | 1,406 | 8.53 | 3,340 | 20.26 | 2.38 |

*The maternity category is calculated using member months for members 10-64 years.

Inpatient Utilization--General Hospital/Acute Care: Dual Eligibles (IPUB)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months |
|--------------|----------------|
| <1 | 23,881 |
| 1-9 | 136,091 |
| 10-19 | 90,186 |
| 20-44 | 59,213 |
| 45-64 | 15,420 |
| 65-74 | 8 |
| 75-84 | 0 |
| 85+ | 0 |
| Unknown | 0 |
| Total | 324,799 |

| Total Inpatient | | | | | |
|-----------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 222 | 9.30 | 1328 | 55.61 | 5.98 |
| 1-9 | 313 | 2.30 | 1044 | 7.67 | 3.34 |
| 10-19 | 384 | 4.26 | 1221 | 13.54 | 3.18 |
| 20-44 | 1485 | 25.08 | 4188 | 70.73 | 2.82 |
| 45-64 | 294 | 19.07 | 1727 | 112.00 | 5.87 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 2,698 | 8.31 | 9,508 | 29.27 | 3.52 |

| Medicine | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 189 | 7.91 | 753 | 31.53 | 3.98 |
| 1-9 | 261 | 1.92 | 817 | 6.00 | 3.13 |
| 10-19 | 118 | 1.31 | 533 | 5.91 | 4.52 |
| 20-44 | 227 | 3.83 | 1022 | 17.26 | 4.50 |
| 45-64 | 214 | 13.88 | 1084 | 70.30 | 5.07 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 1,009 | 3.11 | 4,209 | 12.96 | 4.17 |

| Surgery | | | | | |
|--------------|------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 33 | 1.38 | 575 | 24.08 | 17.42 |
| 1-9 | 52 | 0.38 | 227 | 1.67 | 4.37 |
| 10-19 | 39 | 0.43 | 175 | 1.94 | 4.49 |
| 20-44 | 82 | 1.38 | 349 | 5.89 | 4.26 |
| 45-64 | 77 | 4.99 | 633 | 41.05 | 8.22 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 283 | 0.87 | 1,959 | 6.03 | 6.92 |

| Maternity* | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| 10-19 | 227 | 2.52 | 513 | 5.69 | 2.26 |
| 20-44 | 1176 | 19.86 | 2817 | 47.57 | 2.40 |
| 45-64 | 3 | 0.19 | 10 | 0.65 | 3.33 |
| Unknown | 0 | | 0 | | NA |
| Total | 1,406 | 8.53 | 3,340 | 20.26 | 2.38 |

*The maternity category is calculated using member months for members 10-64 years.

Inpatient Utilization--General Hospital/Acute Care: Disabled (IPUC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months |
|--------------|----------------|
| <1 | 23,881 |
| 1-9 | 136,091 |
| 10-19 | 90,186 |
| 20-44 | 59,213 |
| 45-64 | 15,420 |
| 65-74 | 8 |
| 75-84 | 0 |
| 85+ | 0 |
| Unknown | 0 |
| Total | 324,799 |

| Total Inpatient | | | | | |
|-----------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 222 | 9.30 | 1328 | 55.61 | 5.98 |
| 1-9 | 313 | 2.30 | 1044 | 7.67 | 3.34 |
| 10-19 | 384 | 4.26 | 1221 | 13.54 | 3.18 |
| 20-44 | 1485 | 25.08 | 4188 | 70.73 | 2.82 |
| 45-64 | 294 | 19.07 | 1727 | 112.00 | 5.87 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 2,698 | 8.31 | 9,508 | 29.27 | 3.52 |

| Medicine | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 189 | 7.91 | 753 | 31.53 | 3.98 |
| 1-9 | 261 | 1.92 | 817 | 6.00 | 3.13 |
| 10-19 | 118 | 1.31 | 533 | 5.91 | 4.52 |
| 20-44 | 227 | 3.83 | 1022 | 17.26 | 4.50 |
| 45-64 | 214 | 13.88 | 1084 | 70.30 | 5.07 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 1,009 | 3.11 | 4,209 | 12.96 | 4.17 |

| Surgery | | | | | |
|--------------|------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 33 | 1.38 | 575 | 24.08 | 17.42 |
| 1-9 | 52 | 0.38 | 227 | 1.67 | 4.37 |
| 10-19 | 39 | 0.43 | 175 | 1.94 | 4.49 |
| 20-44 | 82 | 1.38 | 349 | 5.89 | 4.26 |
| 45-64 | 77 | 4.99 | 633 | 41.05 | 8.22 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 283 | 0.87 | 1,959 | 6.03 | 6.92 |

| Maternity* | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| 10-19 | 227 | 2.52 | 513 | 5.69 | 2.26 |
| 20-44 | 1176 | 19.86 | 2817 | 47.57 | 2.40 |
| 45-64 | 3 | 0.19 | 10 | 0.65 | 3.33 |
| Unknown | 0 | | 0 | | NA |
| Total | 1,406 | 8.53 | 3,340 | 20.26 | 2.38 |

*The maternity category is calculated using member months for members 10-64 years.

Inpatient Utilization--General Hospital/Acute Care: Other (IPUD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months |
|--------------|----------------|
| <1 | 23,881 |
| 1-9 | 136,091 |
| 10-19 | 90,186 |
| 20-44 | 59,213 |
| 45-64 | 15,420 |
| 65-74 | 8 |
| 75-84 | 0 |
| 85+ | 0 |
| Unknown | 0 |
| Total | 324,799 |

| Total Inpatient | | | | | |
|-----------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 222 | 9.30 | 1328 | 55.61 | 5.98 |
| 1-9 | 313 | 2.30 | 1044 | 7.67 | 3.34 |
| 10-19 | 384 | 4.26 | 1221 | 13.54 | 3.18 |
| 20-44 | 1485 | 25.08 | 4188 | 70.73 | 2.82 |
| 45-64 | 294 | 19.07 | 1727 | 112.00 | 5.87 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 2,698 | 8.31 | 9,508 | 29.27 | 3.52 |

| Medicine | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 189 | 7.91 | 753 | 31.53 | 3.98 |
| 1-9 | 261 | 1.92 | 817 | 6.00 | 3.13 |
| 10-19 | 118 | 1.31 | 533 | 5.91 | 4.52 |
| 20-44 | 227 | 3.83 | 1022 | 17.26 | 4.50 |
| 45-64 | 214 | 13.88 | 1084 | 70.30 | 5.07 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 1,009 | 3.11 | 4,209 | 12.96 | 4.17 |

| Surgery | | | | | |
|--------------|------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| <1 | 33 | 1.38 | 575 | 24.08 | 17.42 |
| 1-9 | 52 | 0.38 | 227 | 1.67 | 4.37 |
| 10-19 | 39 | 0.43 | 175 | 1.94 | 4.49 |
| 20-44 | 82 | 1.38 | 349 | 5.89 | 4.26 |
| 45-64 | 77 | 4.99 | 633 | 41.05 | 8.22 |
| 65-74 | 0 | 0.00 | 0 | 0.00 | NA |
| 75-84 | 0 | NA | 0 | NA | NA |
| 85+ | 0 | NA | 0 | NA | NA |
| Unknown | 0 | | 0 | | NA |
| Total | 283 | 0.87 | 1,959 | 6.03 | 6.92 |

| Maternity* | | | | | |
|--------------|--------------|----------------------------------|--------------|-----------------------------|------------------------|
| Age | Discharges | Discharges / 1,000 Member Months | Days | Days / 1,000 Members Months | Average Length of Stay |
| 10-19 | 227 | 2.52 | 513 | 5.69 | 2.26 |
| 20-44 | 1176 | 19.86 | 2817 | 47.57 | 2.40 |
| 45-64 | 3 | 0.19 | 10 | 0.65 | 3.33 |
| Unknown | 0 | | 0 | | NA |
| Total | 1,406 | 8.53 | 3,340 | 20.26 | 2.38 |

*The maternity category is calculated using member months for members 10-64 years.

Identification of Alcohol and Other Drug Services: Total (IADA)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|---------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| | | | | | | | | | | | | |
|--------------|----|----|----|----|----|----|----|----|----|----|----|----|
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
|--------------|----|----|----|----|----|----|----|----|----|----|----|----|

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| | | 0-12 | M | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Identification of Alcohol and Other Drug Services: Dual Eligibles (IADB)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Identification of Alcohol and Other Drug Services: Disabled (IADC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Identification of Alcohol and Other Drug Services: Other (IADD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-24 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 25-34 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 35-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Mental Health Utilization: Total (MPTA)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Mental Health Utilization: Dual Eligibles (MPTB)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Mental Health Utilization: Disabled (MPTC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Mental Health Utilization: Other (MPTD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Member Months (Any) | | | Member Months (Inpatient) | | | Member Months (Intensive Outpatient/Partial Hospitalization) | | | Member Months (Outpatient/ED) | | |
|--------------|---------------------|--------|-------|---------------------------|--------|-------|--|--------|-------|-------------------------------|--------|-------|
| | Male | Female | Total | Male | Female | Total | Male | Female | Total | Male | Female | Total |
| 0-12 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Age | Sex | Any Services | | Inpatient | | Intensive Outpatient/Partial Hospitalization | | Outpatient/ED | |
|--------------|--------------|--------------|---------|-----------|---------|--|---------|---------------|---------|
| | | Number | Percent | Number | Percent | Number | Percent | Number | Percent |
| 0-12 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 13-17 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 18-64 | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| 65+ | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | M | NR | NR | NR | NR | NR | NR | NR | NR |
| | F | NR | NR | NR | NR | NR | NR | NR | NR |
| | Total | NR | NR | NR | NR | NR | NR | NR | NR |

Antibiotic Utilization: Total (ABXA)


Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Pharmacy Benefit Member Months | | | |
|--------------------------------|----------------|----------------|----------------|
| Age | Male | Female | Total |
| 0-9 | 82181 | 77791 | 159,972 |
| 10-17 | 40309 | 39180 | 79,489 |
| 18-34 | 15260 | 39976 | 55,236 |
| 35-49 | 7270 | 12144 | 19,414 |
| 50-64 | 5939 | 4741 | 10,680 |
| 65-74 | 3 | 5 | 8 |
| 75-84 | 0 | 0 | 0 |
| 85+ | 0 | 0 | 0 |
| Unknown | 0 | 0 | 0 |
| Total | 150,962 | 173,837 | 324,799 |

| Antibiotic Utilization | | | | | | | | |
|------------------------|--------------|--------------------------|--------------------------------------|--|--|--|---|--|
| Age | Sex | Total Antibiotic Scripts | Average Scripts PMPY for Antibiotics | Total Days Supplied for All Antibiotic Scripts | Average Days Supplied per Antibiotic Scrip | Total Number of Scripts for Antibiotics of Concern | Average Scripts PMPY for Antibiotics of Concern | Percentage of Antibiotics of Concern of all Antibiotic Scripts |
| 0-9 | M | 6582 | 0.96 | 57957 | 8.81 | 2900 | 0.42 | 44.06% |
| | F | 6139 | 0.95 | 54203 | 8.83 | 2519 | 0.39 | 41.03% |
| | Total | 12,721 | 0.95 | 112,160 | 8.82 | 5,419 | 0.41 | 42.60% |
| 10-17 | M | 1880 | 0.56 | 18789 | 9.99 | 866 | 0.26 | 46.06% |
| | F | 2480 | 0.76 | 22412 | 9.04 | 1058 | 0.32 | 42.66% |
| | Total | 4,360 | 0.66 | 41,201 | 9.45 | 1,924 | 0.29 | 44.13% |
| 18-34 | M | 958 | 0.75 | 9030 | 9.43 | 413 | 0.32 | 43.11% |
| | F | 4861 | 1.46 | 39759 | 8.18 | 1912 | 0.57 | 39.33% |
| | Total | 5,819 | 1.26 | 48,789 | 8.38 | 2,325 | 0.51 | 39.96% |
| 35-49 | M | 662 | 1.09 | 7142 | 10.79 | 306 | 0.51 | 46.22% |
| | F | 1651 | 1.63 | 14365 | 8.70 | 813 | 0.80 | 49.24% |
| | Total | 2,313 | 1.43 | 21,507 | 9.30 | 1,119 | 0.69 | 48.38% |
| 50-64 | M | 546 | 1.10 | 5631 | 10.31 | 220 | 0.44 | 40.29% |
| | F | 617 | 1.56 | 5261 | 8.53 | 280 | 0.71 | 45.38% |
| | Total | 1,163 | 1.31 | 10,892 | 9.37 | 500 | 0.56 | 42.99% |
| 65-74 | M | 0 | 0.00 | 0 | NA | 0 | 0.00 | NA |
| | F | 1 | 2.40 | 30 | 30.00 | 0 | 0.00 | 0.00% |
| | Total | 1 | 1.50 | 30 | 30.00 | 0 | 0.00 | 0.00% |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Total | M | 10,628 | 0.84 | 98,549 | 9.27 | 4,705 | 0.37 | 44.27% |
| | F | 15,749 | 1.09 | 136,030 | 8.64 | 6,582 | 0.45 | 41.79% |
| | Total | 26,377 | 0.97 | 234,579 | 8.89 | 11,287 | 0.42 | 42.79% |

| Antibiotics of Concern Utilization | | | | | | | | |
|------------------------------------|-------|-------------------------|-------------------------------------|--|--|---|--|---------------------------------------|
| Age | Sex | Total Quinolone Scripts | Average Scripts PMPY for Quinolones | Total Cephalosporin 2nd-4th Generation Scripts | Average Scripts PMPY for Cephalosporins 2nd-4th Generation | Total Azithromycin and Clarithromycin Scripts | Average Scripts PMPY for Azithromycin and Clarithromycin | Total Amoxicillin/Clavulanate Scripts |
| 0-9 | M | 2 | 0.00 | 422 | 0.06 | 1673 | 0.24 | 769 |
| | F | 3 | 0.00 | 350 | 0.05 | 1408 | 0.22 | 708 |
| | Total | 5 | 0.00 | 772 | 0.06 | 3,081 | 0.23 | 1,477 |
| 10-17 | M | 20 | 0.01 | 45 | 0.01 | 561 | 0.17 | 206 |
| | F | 64 | 0.02 | 75 | 0.02 | 674 | 0.21 | 204 |
| | Total | 84 | 0.01 | 120 | 0.02 | 1,235 | 0.19 | 410 |
| 18-34 | M | 59 | 0.05 | 16 | 0.01 | 210 | 0.17 | 80 |
| | F | 362 | 0.11 | 66 | 0.02 | 942 | 0.28 | 330 |
| | Total | 421 | 0.09 | 82 | 0.02 | 1,152 | 0.25 | 410 |
| 35-49 | M | 86 | 0.14 | 16 | 0.03 | 106 | 0.17 | 58 |
| | F | 191 | 0.19 | 26 | 0.03 | 387 | 0.38 | 118 |
| | Total | 277 | 0.17 | 42 | 0.03 | 493 | 0.30 | 176 |
| 50-64 | M | 55 | 0.11 | 5 | 0.01 | 83 | 0.17 | 42 |
| | F | 82 | 0.21 | 3 | 0.01 | 128 | 0.32 | 38 |
| | Total | 137 | 0.15 | 8 | 0.01 | 211 | 0.24 | 80 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| Total | M | 222 | 0.02 | 504 | 0.04 | 2,633 | 0.21 | 1,155 |
| | F | 702 | 0.05 | 520 | 0.04 | 3,539 | 0.24 | 1,398 |
| | Total | 924 | 0.03 | 1,024 | 0.04 | 6,172 | 0.23 | 2,553 |

| All Other Antibiotics Utilization | | | | | | | | |
|-----------------------------------|-------|--------------------------------------|--|------------------------------|--|--|--|---------------------------|
| Age | Sex | Total Absorbable Sulfonamide Scripts | Average Scripts PMPY for Absorbable Sulfonamides | Total Aminoglycoside Scripts | Average Scripts PMPY for Aminoglycosides | Total 1st Generation Cephalosporin Scripts | Average Scripts PMPY for 1st Generation Cephalosporins | Total Lincosamide Scripts |
| 0-9 | M | 321 | 0.05 | 1 | 0.00 | 405 | 0.06 | 0 |
| | F | 382 | 0.06 | 0 | 0.00 | 500 | 0.08 | 0 |
| | Total | 703 | 0.05 | 1 | 0.00 | 905 | 0.07 | 0 |
| 10-17 | M | 84 | 0.03 | 0 | 0.00 | 194 | 0.06 | 0 |
| | F | 196 | 0.06 | 3 | 0.00 | 301 | 0.09 | 0 |
| | Total | 280 | 0.04 | 3 | 0.00 | 495 | 0.07 | 0 |
| 18-34 | M | 85 | 0.07 | 0 | 0.00 | 92 | 0.07 | 0 |
| | F | 372 | 0.11 | 0 | 0.00 | 502 | 0.15 | 0 |
| | Total | 457 | 0.10 | 0 | 0.00 | 594 | 0.13 | 0 |
| 35-49 | M | 65 | 0.11 | 0 | 0.00 | 59 | 0.10 | 0 |
| | F | 144 | 0.14 | 0 | 0.00 | 153 | 0.15 | 0 |
| | Total | 209 | 0.13 | 0 | 0.00 | 212 | 0.13 | 0 |
| 50-64 | M | 98 | 0.20 | 0 | 0.00 | 51 | 0.10 | 0 |
| | F | 75 | 0.19 | 0 | 0.00 | 64 | 0.16 | 0 |
| | Total | 173 | 0.19 | 0 | 0.00 | 115 | 0.13 | 0 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 |
| Total | M | 653 | 0.05 | 1 | 0.00 | 801 | 0.06 | 0 |
| | F | 1,169 | 0.08 | 3 | 0.00 | 1,520 | 0.10 | 0 |
| | Total | 1,822 | 0.07 | 4 | 0.00 | 2,321 | 0.09 | 0 |



| Average Scrips PMPY for Amoxicillin/Clavulanates | Total Ketolides Scrips | Average Scrips PMPY for Ketolides | Total Clindamycin Scrips | Average Scrips PMPY for Clindamycins | Total Misc. Antibiotics of Concern Scrips | Average Scrips PMPY for Misc. Antibiotics of Concern |
|--|------------------------|-----------------------------------|--------------------------|--------------------------------------|---|--|
| 0.11 | 0 | 0.00 | 34 | 0.00 | 0 | 0.00 |
| 0.11 | 0 | 0.00 | 49 | 0.01 | 1 | 0.00 |
| 0.11 | 0 | 0.00 | 83 | 0.01 | 1 | 0.00 |
| 0.06 | 0 | 0.00 | 29 | 0.01 | 5 | 0.00 |
| 0.06 | 0 | 0.00 | 39 | 0.01 | 2 | 0.00 |
| 0.06 | 0 | 0.00 | 68 | 0.01 | 7 | 0.00 |
| 0.06 | 0 | 0.00 | 46 | 0.04 | 2 | 0.00 |
| 0.10 | 0 | 0.00 | 210 | 0.06 | 2 | 0.00 |
| 0.09 | 0 | 0.00 | 256 | 0.06 | 4 | 0.00 |
| 0.10 | 0 | 0.00 | 39 | 0.06 | 1 | 0.00 |
| 0.12 | 0 | 0.00 | 89 | 0.09 | 2 | 0.00 |
| 0.11 | 0 | 0.00 | 128 | 0.08 | 3 | 0.00 |
| 0.08 | 0 | 0.00 | 33 | 0.07 | 2 | 0.00 |
| 0.10 | 0 | 0.00 | 28 | 0.07 | 1 | 0.00 |
| 0.09 | 0 | 0.00 | 61 | 0.07 | 3 | 0.00 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA |
| 0.09 | 0 | 0.00 | 181 | 0.01 | 10 | 0.00 |
| 0.10 | 0 | 0.00 | 415 | 0.03 | 8 | 0.00 |
| 0.09 | 0 | 0.00 | 596 | 0.02 | 18 | 0.00 |

on

| Average Scrips PMPY for Lincosamides | Total Macrolides (not azith. or clarith.) Scrips | Average Scrips PMPY for Macrolides (not azith. or clarith.) | Total Penicillin Scrips | Average Scrips PMPY for Penicillins | Total Tetracycline Scrips | Average Scrips PMPY for Tetracyclines | Total Misc. Antibiotic Scrips | Average Scrips PMPY for Misc. Antibiotics |
|--------------------------------------|--|---|-------------------------|-------------------------------------|---------------------------|---------------------------------------|-------------------------------|---|
| 0.00 | 14 | 0.00 | 2919 | 0.43 | 0 | 0.00 | 22 | 0.00 |
| 0.00 | 6 | 0.00 | 2669 | 0.41 | 7 | 0.00 | 56 | 0.01 |
| 0.00 | 20 | 0.00 | 5,588 | 0.42 | 7 | 0.00 | 78 | 0.01 |
| 0.00 | 5 | 0.00 | 566 | 0.17 | 157 | 0.05 | 8 | 0.00 |
| 0.00 | 11 | 0.00 | 652 | 0.20 | 132 | 0.04 | 127 | 0.04 |
| 0.00 | 16 | 0.00 | 1,218 | 0.18 | 289 | 0.04 | 135 | 0.02 |
| 0.00 | 5 | 0.00 | 256 | 0.20 | 88 | 0.07 | 19 | 0.01 |
| 0.00 | 11 | 0.00 | 920 | 0.28 | 256 | 0.08 | 888 | 0.27 |
| 0.00 | 16 | 0.00 | 1,176 | 0.26 | 344 | 0.07 | 907 | 0.20 |
| 0.00 | 6 | 0.01 | 176 | 0.29 | 34 | 0.06 | 16 | 0.03 |
| 0.00 | 1 | 0.00 | 275 | 0.27 | 91 | 0.09 | 174 | 0.17 |
| 0.00 | 7 | 0.00 | 451 | 0.28 | 125 | 0.08 | 190 | 0.12 |
| 0.00 | 0 | 0.00 | 124 | 0.25 | 38 | 0.08 | 15 | 0.03 |
| 0.00 | 0 | 0.00 | 104 | 0.26 | 29 | 0.07 | 65 | 0.16 |
| 0.00 | 0 | 0.00 | 228 | 0.26 | 67 | 0.08 | 80 | 0.09 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 2.40 |
| 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 1.50 |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 0.00 | 30 | 0.00 | 4,041 | 0.32 | 317 | 0.03 | 80 | 0.01 |
| 0.00 | 29 | 0.00 | 4,620 | 0.32 | 515 | 0.04 | 1,311 | 0.09 |
| 0.00 | 59 | 0.00 | 8,661 | 0.32 | 832 | 0.03 | 1,391 | 0.05 |

Antibiotic Utilization: Dual Eligibles (ABXB)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Member Months | | | |
|---------------|----------------|----------------|----------------|
| Age | Member Months | | |
| | Male | Female | Total |
| 0-9 | 82181 | 77791 | 159,972 |
| 10-17 | 40309 | 39180 | 79,489 |
| 18-34 | 15260 | 39976 | 55,236 |
| 35-49 | 7270 | 12144 | 19,414 |
| 50-64 | 5939 | 4741 | 10,680 |
| 65-74 | 3 | 5 | 8 |
| 75-84 | 0 | 0 | 0 |
| 85+ | 0 | 0 | 0 |
| Unknown | 0 | 0 | 0 |
| Total | 150,962 | 173,837 | 324,799 |

| Antibiotic Utilization | | | | | | | | |
|------------------------|--------------|--------------------------|--------------------------------------|--|--|--|---|--|
| Age | Sex | Total Antibiotic Scripts | Average Scripts PMPY for Antibiotics | Total Days Supplied for All Antibiotic Scripts | Average Days Supplied per Antibiotic Scrip | Total Number of Scripts for Antibiotics of Concern | Average Scripts PMPY for Antibiotics of Concern | Percentage of Antibiotics of Concern of all Antibiotic Scripts |
| 0-9 | M | 6582 | 0.96 | 57957 | 8.81 | 2900 | 0.42 | 44% |
| | F | 6139 | 0.95 | 54203 | 8.83 | 2519 | 0.39 | 41% |
| | Total | 12,721 | 0.95 | 112,160 | 8.82 | 5,419 | 0.41 | 43% |
| 10-17 | M | 1880 | 0.56 | 18789 | 9.99 | 866 | 0.26 | 46% |
| | F | 2480 | 0.76 | 22412 | 9.04 | 1058 | 0.32 | 43% |
| | Total | 4,360 | 0.66 | 41,201 | 9.45 | 1,924 | 0.29 | 44% |
| 18-34 | M | 958 | 0.75 | 9030 | 9.43 | 413 | 0.32 | 43% |
| | F | 4861 | 1.46 | 39759 | 8.18 | 1912 | 0.57 | 39% |
| | Total | 5,819 | 1.26 | 48,789 | 8.38 | 2,325 | 0.51 | 40% |
| 35-49 | M | 662 | 1.09 | 7142 | 10.79 | 306 | 0.51 | 46% |
| | F | 1651 | 1.63 | 14365 | 8.70 | 813 | 0.80 | 49% |
| | Total | 2,313 | 1.43 | 21,507 | 9.30 | 1,119 | 0.69 | 48% |
| 50-64 | M | 546 | 1.10 | 5631 | 10.31 | 220 | 0.44 | 40% |
| | F | 617 | 1.56 | 5261 | 8.53 | 280 | 0.71 | 45% |
| | Total | 1,163 | 1.31 | 10,892 | 9.37 | 500 | 0.56 | 43% |
| 65-74 | M | 0 | 0.00 | 0 | NA | 0 | 0.00 | NA |
| | F | 1 | 2.40 | 30 | 30.00 | 0 | 0.00 | 0% |
| | Total | 1 | 1.50 | 30 | 30.00 | 0 | 0.00 | 0% |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Total | M | 10,628 | 0.84 | 98,549 | 9.27 | 4,705 | 0.37 | 44% |
| | F | 15,749 | 1.09 | 136,030 | 8.64 | 6,582 | 0.45 | 42% |
| | Total | 26,377 | 0.97 | 234,579 | 8.89 | 11,287 | 0.42 | 43% |

| Antibiotics of Concern Utilization | | | | | | | | | | | | | | | |
|------------------------------------|--------------|-------------------------|-------------------------------------|--|--|---|--|---------------------------------------|---|-------------------------|------------------------------------|---------------------------|---------------------------------------|--|---|
| Age | Sex | Total Quinolone Scripts | Average Scripts PMPY for Quinolones | Total Cephalosporin 2nd-4th Generation Scripts | Average Scripts PMPY for Cephalosporins 2nd-4th Generation | Total Azithromycin and Clarithromycin Scripts | Average Scripts PMPY for Azithromycins and Clarithromycins | Total Amoxicillin/Clavulanate Scripts | Average Scripts PMPY for Amoxicillin/Clavulanates | Total Ketolides Scripts | Average Scripts PMPY for Ketolides | Total Clindamycin Scripts | Average Scripts PMPY for Clindamycins | Total Misc. Antibiotics of Concern Scripts | Average Scripts PMPY for Misc. Antibiotics of Concern |
| 0-9 | M | 2 | 0.00 | 422 | 0.06 | 1673 | 0.24 | 769 | 0.11 | 0 | 0.00 | 34 | 0.00 | 0 | 0.00 |
| | F | 3 | 0.00 | 350 | 0.05 | 1408 | 0.22 | 708 | 0.11 | 0 | 0.00 | 49 | 0.01 | 1 | 0.00 |
| | Total | 5 | 0.00 | 772 | 0.06 | 3,081 | 0.23 | 1,477 | 0.11 | 0 | 0.00 | 83 | 0.01 | 1 | 0.00 |
| 10-17 | M | 20 | 0.01 | 45 | 0.01 | 561 | 0.17 | 206 | 0.06 | 0 | 0.00 | 29 | 0.01 | 5 | 0.00 |
| | F | 64 | 0.02 | 75 | 0.02 | 674 | 0.21 | 204 | 0.06 | 0 | 0.00 | 39 | 0.01 | 2 | 0.00 |
| | Total | 84 | 0.01 | 120 | 0.02 | 1,235 | 0.19 | 410 | 0.06 | 0 | 0.00 | 68 | 0.01 | 7 | 0.00 |
| 18-34 | M | 59 | 0.05 | 16 | 0.01 | 210 | 0.17 | 80 | 0.06 | 0 | 0.00 | 46 | 0.04 | 2 | 0.00 |
| | F | 362 | 0.11 | 66 | 0.02 | 942 | 0.28 | 330 | 0.10 | 0 | 0.00 | 210 | 0.06 | 2 | 0.00 |
| | Total | 421 | 0.09 | 82 | 0.02 | 1,152 | 0.25 | 410 | 0.09 | 0 | 0.00 | 256 | 0.06 | 4 | 0.00 |
| 35-49 | M | 86 | 0.14 | 16 | 0.03 | 106 | 0.17 | 58 | 0.10 | 0 | 0.00 | 39 | 0.06 | 1 | 0.00 |
| | F | 191 | 0.19 | 26 | 0.03 | 387 | 0.38 | 118 | 0.12 | 0 | 0.00 | 89 | 0.09 | 2 | 0.00 |
| | Total | 277 | 0.17 | 42 | 0.03 | 493 | 0.30 | 176 | 0.11 | 0 | 0.00 | 128 | 0.08 | 3 | 0.00 |
| 50-64 | M | 55 | 0.11 | 5 | 0.01 | 83 | 0.17 | 42 | 0.08 | 0 | 0.00 | 33 | 0.07 | 2 | 0.00 |
| | F | 82 | 0.21 | 3 | 0.01 | 128 | 0.32 | 38 | 0.10 | 0 | 0.00 | 28 | 0.07 | 1 | 0.00 |
| | Total | 137 | 0.15 | 8 | 0.01 | 211 | 0.24 | 80 | 0.09 | 0 | 0.00 | 61 | 0.07 | 3 | 0.00 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 222 | 0.02 | 504 | 0.04 | 2,633 | 0.21 | 1,155 | 0.09 | 0 | 0.00 | 181 | 0.01 | 10 | 0.00 |
| | F | 702 | 0.05 | 520 | 0.04 | 3,539 | 0.24 | 1,398 | 0.10 | 0 | 0.00 | 415 | 0.03 | 8 | 0.00 |
| | Total | 924 | 0.03 | 1,024 | 0.04 | 6,172 | 0.23 | 2,553 | 0.09 | 0 | 0.00 | 596 | 0.02 | 18 | 0.00 |

| All Other Antibiotics Utilization | | | | | | | | | | | | | | | | | |
|-----------------------------------|-------|-------------------------------------|---|-----------------------------|---|---|---|--------------------------|--------------------------------------|--|---|-------------------------|-------------------------------------|---------------------------|---------------------------------------|-------------------------------|---|
| Age | Sex | Total Absorbable Sulfonamide Scrips | Average Scrips PMPY for Absorbable Sulfonamides | Total Aminoglycoside Scrips | Average Scrips PMPY for Aminoglycosides | Total 1st Generation Cephalosporin Scrips | Average Scrips PMPY for 1st Generation Cephalosporins | Total Lincosamide Scrips | Average Scrips PMPY for Lincosamides | Total Macrolides (not azith. or clarith.) Scrips | Average Scrips PMPY for Macrolides (not azith. or clarith.) | Total Penicillin Scrips | Average Scrips PMPY for Penicillins | Total Tetracycline Scrips | Average Scrips PMPY for Tetracyclines | Total Misc. Antibiotic Scrips | Average Scrips PMPY for Misc. Antibiotics |
| 0-9 | M | 321 | 0.05 | 1 | 0.00 | 405 | 0.06 | 0 | 0.00 | 14 | 0.00 | 2919 | 0.43 | 0 | 0.00 | 22 | 0.00 |
| | F | 382 | 0.06 | 0 | 0.00 | 500 | 0.08 | 0 | 0.00 | 6 | 0.00 | 2669 | 0.41 | 7 | 0.00 | 56 | 0.01 |
| | Total | 703 | 0.05 | 1 | 0.00 | 905 | 0.07 | 0 | 0.00 | 20 | 0.00 | 5,588 | 0.42 | 7 | 0.00 | 78 | 0.01 |
| 10-17 | M | 84 | 0.03 | 0 | 0.00 | 194 | 0.06 | 0 | 0.00 | 5 | 0.00 | 566 | 0.17 | 157 | 0.05 | 8 | 0.00 |
| | F | 196 | 0.06 | 3 | 0.00 | 301 | 0.09 | 0 | 0.00 | 11 | 0.00 | 652 | 0.20 | 132 | 0.04 | 127 | 0.04 |
| | Total | 280 | 0.04 | 3 | 0.00 | 495 | 0.07 | 0 | 0.00 | 16 | 0.00 | 1,218 | 0.18 | 289 | 0.04 | 135 | 0.02 |
| 18-34 | M | 85 | 0.07 | 0 | 0.00 | 92 | 0.07 | 0 | 0.00 | 5 | 0.00 | 256 | 0.20 | 88 | 0.07 | 19 | 0.01 |
| | F | 372 | 0.11 | 0 | 0.00 | 502 | 0.15 | 0 | 0.00 | 11 | 0.00 | 920 | 0.28 | 256 | 0.08 | 888 | 0.27 |
| | Total | 457 | 0.10 | 0 | 0.00 | 594 | 0.13 | 0 | 0.00 | 16 | 0.00 | 1,176 | 0.26 | 344 | 0.07 | 907 | 0.20 |
| 35-49 | M | 65 | 0.11 | 0 | 0.00 | 59 | 0.10 | 0 | 0.00 | 6 | 0.01 | 176 | 0.29 | 34 | 0.06 | 16 | 0.03 |
| | F | 144 | 0.14 | 0 | 0.00 | 153 | 0.15 | 0 | 0.00 | 1 | 0.00 | 275 | 0.27 | 91 | 0.09 | 174 | 0.17 |
| | Total | 209 | 0.13 | 0 | 0.00 | 212 | 0.13 | 0 | 0.00 | 7 | 0.00 | 451 | 0.28 | 125 | 0.08 | 190 | 0.12 |
| 50-64 | M | 98 | 0.20 | 0 | 0.00 | 51 | 0.10 | 0 | 0.00 | 0 | 0.00 | 124 | 0.25 | 38 | 0.08 | 15 | 0.03 |
| | F | 75 | 0.19 | 0 | 0.00 | 64 | 0.16 | 0 | 0.00 | 0 | 0.00 | 104 | 0.26 | 29 | 0.07 | 65 | 0.16 |
| | Total | 173 | 0.19 | 0 | 0.00 | 115 | 0.13 | 0 | 0.00 | 0 | 0.00 | 228 | 0.26 | 67 | 0.08 | 80 | 0.09 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 2.40 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 1.50 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 653 | 0.05 | 1 | 0.00 | 801 | 0.06 | 0 | 0.00 | 30 | 0.00 | 4,041 | 0.32 | 317 | 0.03 | 80 | 0.01 |
| | F | 1,169 | 0.08 | 3 | 0.00 | 1,520 | 0.10 | 0 | 0.00 | 29 | 0.00 | 4,620 | 0.32 | 515 | 0.04 | 1,311 | 0.09 |
| | Total | 1,822 | 0.07 | 4 | 0.00 | 2,321 | 0.09 | 0 | 0.00 | 59 | 0.00 | 8,661 | 0.32 | 832 | 0.03 | 1,391 | 0.05 |

Antibiotic Utilization: Disabled (ABXC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Member Months | | | |
|---------------|----------------|----------------|----------------|
| Age | Member Months | | |
| | Male | Female | Total |
| 0-9 | 82181 | 77791 | 159,972 |
| 10-17 | 40309 | 39180 | 79,489 |
| 18-34 | 15260 | 39976 | 55,236 |
| 35-49 | 7270 | 12144 | 19,414 |
| 50-64 | 5939 | 4741 | 10,680 |
| 65-74 | 3 | 5 | 8 |
| 75-84 | 0 | 0 | 0 |
| 85+ | 0 | 0 | 0 |
| Unknown | 0 | 0 | 0 |
| Total | 150,962 | 173,837 | 324,799 |

| Antibiotic Utilization | | | | | | | | |
|------------------------|--------------|--------------------------|--------------------------------------|--|--|--|---|--|
| Age | Sex | Total Antibiotic Scripts | Average Scripts PMPY for Antibiotics | Total Days Supplied for All Antibiotic Scripts | Average Days Supplied per Antibiotic Scrip | Total Number of Scripts for Antibiotics of Concern | Average Scripts PMPY for Antibiotics of Concern | Percentage of Antibiotics of Concern of all Antibiotic Scripts |
| 0-9 | M | 6582 | 0.96 | 57957 | 8.81 | 2900 | 0.42 | 44% |
| | F | 6139 | 0.95 | 54203 | 8.83 | 2519 | 0.39 | 41% |
| | Total | 12,721 | 0.95 | 112,160 | 8.82 | 5,419 | 0.41 | 43% |
| 10-17 | M | 1880 | 0.56 | 18789 | 9.99 | 866 | 0.26 | 46% |
| | F | 2480 | 0.76 | 22412 | 9.04 | 1058 | 0.32 | 43% |
| | Total | 4,360 | 0.66 | 41,201 | 9.45 | 1,924 | 0.29 | 44% |
| 18-34 | M | 958 | 0.75 | 9030 | 9.43 | 413 | 0.32 | 43% |
| | F | 4861 | 1.46 | 39759 | 8.18 | 1912 | 0.57 | 39% |
| | Total | 5,819 | 1.26 | 48,789 | 8.38 | 2,325 | 0.51 | 40% |

| | | | | | | | | |
|---------|-------|--------|------|---------|-------|--------|------|-----|
| 35-49 | M | 662 | 1.09 | 7142 | 10.79 | 306 | 0.51 | 46% |
| | F | 1651 | 1.63 | 14365 | 8.70 | 813 | 0.80 | 49% |
| | Total | 2,313 | 1.43 | 21,507 | 9.30 | 1,119 | 0.69 | 48% |
| 50-64 | M | 546 | 1.10 | 5631 | 10.31 | 220 | 0.44 | 40% |
| | F | 617 | 1.56 | 5261 | 8.53 | 280 | 0.71 | 45% |
| | Total | 1,163 | 1.31 | 10,892 | 9.37 | 500 | 0.56 | 43% |
| 65-74 | M | 0 | 0.00 | 0 | NA | 0 | 0.00 | NA |
| | F | 1 | 2.40 | 30 | 30.00 | 0 | 0.00 | 0% |
| | Total | 1 | 1.50 | 30 | 30.00 | 0 | 0.00 | 0% |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Total | M | 10,628 | 0.84 | 98,549 | 9.27 | 4,705 | 0.37 | 44% |
| | F | 15,749 | 1.09 | 136,030 | 8.64 | 6,582 | 0.45 | 42% |
| | Total | 26,377 | 0.97 | 234,579 | 8.89 | 11,287 | 0.42 | 43% |

| Antibiotics of Concern Utilization | | | | | | | | | | | | | | | |
|------------------------------------|--------------|-------------------------|-------------------------------------|--|--|---|--|---------------------------------------|---|-------------------------|------------------------------------|---------------------------|---------------------------------------|--|---|
| Age | Sex | Total Quinolone Scripts | Average Scripts PMPY for Quinolones | Total Cephalosporin 2nd-4th Generation Scripts | Average Scripts PMPY for Cephalosporins 2nd-4th Generation | Total Azithromycin and Clarithromycin Scripts | Average Scripts PMPY for Azithromycins and Clarithromycins | Total Amoxicillin/Clavulanate Scripts | Average Scripts PMPY for Amoxicillin/Clavulanates | Total Ketolides Scripts | Average Scripts PMPY for Ketolides | Total Clindamycin Scripts | Average Scripts PMPY for Clindamycins | Total Misc. Antibiotics of Concern Scripts | Average Scripts PMPY for Misc. Antibiotics of Concern |
| 0-9 | M | 2 | 0.00 | 422 | 0.06 | 1673 | 0.24 | 769 | 0.11 | 0 | 0.00 | 34 | 0.00 | 0 | 0.00 |
| | F | 3 | 0.00 | 350 | 0.05 | 1408 | 0.22 | 708 | 0.11 | 0 | 0.00 | 49 | 0.01 | 1 | 0.00 |
| | Total | 5 | 0.00 | 772 | 0.06 | 3,081 | 0.23 | 1,477 | 0.11 | 0 | 0.00 | 83 | 0.01 | 1 | 0.00 |
| 10-17 | M | 20 | 0.01 | 45 | 0.01 | 561 | 0.17 | 206 | 0.06 | 0 | 0.00 | 29 | 0.01 | 5 | 0.00 |
| | F | 64 | 0.02 | 75 | 0.02 | 674 | 0.21 | 204 | 0.06 | 0 | 0.00 | 39 | 0.01 | 2 | 0.00 |
| | Total | 84 | 0.01 | 120 | 0.02 | 1,235 | 0.19 | 410 | 0.06 | 0 | 0.00 | 68 | 0.01 | 7 | 0.00 |
| 18-34 | M | 59 | 0.05 | 16 | 0.01 | 210 | 0.17 | 80 | 0.06 | 0 | 0.00 | 46 | 0.04 | 2 | 0.00 |
| | F | 362 | 0.11 | 66 | 0.02 | 942 | 0.28 | 330 | 0.10 | 0 | 0.00 | 210 | 0.06 | 2 | 0.00 |
| | Total | 421 | 0.09 | 82 | 0.02 | 1,152 | 0.25 | 410 | 0.09 | 0 | 0.00 | 256 | 0.06 | 4 | 0.00 |
| 35-49 | M | 86 | 0.14 | 16 | 0.03 | 106 | 0.17 | 58 | 0.10 | 0 | 0.00 | 39 | 0.06 | 1 | 0.00 |
| | F | 191 | 0.19 | 26 | 0.03 | 387 | 0.38 | 118 | 0.12 | 0 | 0.00 | 89 | 0.09 | 2 | 0.00 |
| | Total | 277 | 0.17 | 42 | 0.03 | 493 | 0.30 | 176 | 0.11 | 0 | 0.00 | 128 | 0.08 | 3 | 0.00 |
| 50-64 | M | 55 | 0.11 | 5 | 0.01 | 83 | 0.17 | 42 | 0.08 | 0 | 0.00 | 33 | 0.07 | 2 | 0.00 |
| | F | 82 | 0.21 | 3 | 0.01 | 128 | 0.32 | 38 | 0.10 | 0 | 0.00 | 28 | 0.07 | 1 | 0.00 |
| | Total | 137 | 0.15 | 8 | 0.01 | 211 | 0.24 | 80 | 0.09 | 0 | 0.00 | 61 | 0.07 | 3 | 0.00 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 222 | 0.02 | 504 | 0.04 | 2,633 | 0.21 | 1,155 | 0.09 | 0 | 0.00 | 181 | 0.01 | 10 | 0.00 |
| | F | 702 | 0.05 | 520 | 0.04 | 3,539 | 0.24 | 1,398 | 0.10 | 0 | 0.00 | 415 | 0.03 | 8 | 0.00 |
| | Total | 924 | 0.03 | 1,024 | 0.04 | 6,172 | 0.23 | 2,553 | 0.09 | 0 | 0.00 | 596 | 0.02 | 18 | 0.00 |

| All Other Antibiotics Utilization | | | | | | | | | | | | | | | | | |
|-----------------------------------|-------|--------------------------------------|--|------------------------------|--|--|--|---------------------------|---------------------------------------|---|--|--------------------------|--------------------------------------|----------------------------|--|--------------------------------|--|
| Age | Sex | Total Absorbable Sulfonamide Scrrips | Average Scrrips PMPY for Absorbable Sulfonamides | Total Aminoglycoside Scrrips | Average Scrrips PMPY for Aminoglycosides | Total 1st Generation Cephalosporin Scrrips | Average Scrrips PMPY for 1st Generation Cephalosporins | Total Lincosamide Scrrips | Average Scrrips PMPY for Lincosamides | Total Macrolides (not azith. or clarith.) Scrrips | Average Scrrips PMPY for Macrolides (not azith. or clarith.) | Total Penicillin Scrrips | Average Scrrips PMPY for Penicillins | Total Tetracycline Scrrips | Average Scrrips PMPY for Tetracyclines | Total Misc. Antibiotic Scrrips | Average Scrrips PMPY for Misc. Antibiotics |
| 0-9 | M | 321 | 0.05 | 1 | 0.00 | 405 | 0.06 | 0 | 0.00 | 14 | 0.00 | 2919 | 0.43 | 0 | 0.00 | 22 | 0.00 |
| | F | 382 | 0.06 | 0 | 0.00 | 500 | 0.08 | 0 | 0.00 | 6 | 0.00 | 2669 | 0.41 | 7 | 0.00 | 56 | 0.01 |
| | Total | 703 | 0.05 | 1 | 0.00 | 905 | 0.07 | 0 | 0.00 | 20 | 0.00 | 5,588 | 0.42 | 7 | 0.00 | 78 | 0.01 |
| 10-17 | M | 84 | 0.03 | 0 | 0.00 | 194 | 0.06 | 0 | 0.00 | 5 | 0.00 | 566 | 0.17 | 157 | 0.05 | 8 | 0.00 |
| | F | 196 | 0.06 | 3 | 0.00 | 301 | 0.09 | 0 | 0.00 | 11 | 0.00 | 652 | 0.20 | 132 | 0.04 | 127 | 0.04 |
| | Total | 280 | 0.04 | 3 | 0.00 | 495 | 0.07 | 0 | 0.00 | 16 | 0.00 | 1,218 | 0.18 | 289 | 0.04 | 135 | 0.02 |
| 18-34 | M | 85 | 0.07 | 0 | 0.00 | 92 | 0.07 | 0 | 0.00 | 5 | 0.00 | 256 | 0.20 | 88 | 0.07 | 19 | 0.01 |
| | F | 372 | 0.11 | 0 | 0.00 | 502 | 0.15 | 0 | 0.00 | 11 | 0.00 | 920 | 0.28 | 256 | 0.08 | 888 | 0.27 |
| | Total | 457 | 0.10 | 0 | 0.00 | 594 | 0.13 | 0 | 0.00 | 16 | 0.00 | 1,176 | 0.26 | 344 | 0.07 | 907 | 0.20 |
| 35-49 | M | 65 | 0.11 | 0 | 0.00 | 59 | 0.10 | 0 | 0.00 | 6 | 0.01 | 176 | 0.29 | 34 | 0.06 | 16 | 0.03 |
| | F | 144 | 0.14 | 0 | 0.00 | 153 | 0.15 | 0 | 0.00 | 1 | 0.00 | 275 | 0.27 | 91 | 0.09 | 174 | 0.17 |
| | Total | 209 | 0.13 | 0 | 0.00 | 212 | 0.13 | 0 | 0.00 | 7 | 0.00 | 451 | 0.28 | 125 | 0.08 | 190 | 0.12 |
| 50-64 | M | 98 | 0.20 | 0 | 0.00 | 51 | 0.10 | 0 | 0.00 | 0 | 0.00 | 124 | 0.25 | 38 | 0.08 | 15 | 0.03 |
| | F | 75 | 0.19 | 0 | 0.00 | 64 | 0.16 | 0 | 0.00 | 0 | 0.00 | 104 | 0.26 | 29 | 0.07 | 65 | 0.16 |
| | Total | 173 | 0.19 | 0 | 0.00 | 115 | 0.13 | 0 | 0.00 | 0 | 0.00 | 228 | 0.26 | 67 | 0.08 | 80 | 0.09 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 2.40 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 1.50 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 653 | 0.05 | 1 | 0.00 | 801 | 0.06 | 0 | 0.00 | 30 | 0.00 | 4,041 | 0.32 | 317 | 0.03 | 80 | 0.01 |
| | F | 1,169 | 0.08 | 3 | 0.00 | 1,520 | 0.10 | 0 | 0.00 | 29 | 0.00 | 4,620 | 0.32 | 515 | 0.04 | 1,311 | 0.09 |
| | Total | 1,822 | 0.07 | 4 | 0.00 | 2,321 | 0.09 | 0 | 0.00 | 59 | 0.00 | 8,661 | 0.32 | 832 | 0.03 | 1,391 | 0.05 |

Antibiotic Utilization: Other (ABXD)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Member Months | | | |
|---------------|----------------|----------------|----------------|
| Age | Member Months | | |
| | Male | Female | Total |
| 0-9 | 82181 | 77791 | 159,972 |
| 10-17 | 40309 | 39180 | 79,489 |
| 18-34 | 15260 | 39976 | 55,236 |
| 35-49 | 7270 | 12144 | 19,414 |
| 50-64 | 5939 | 4741 | 10,680 |
| 65-74 | 3 | 5 | 8 |
| 75-84 | 0 | 0 | 0 |
| 85+ | 0 | 0 | 0 |
| Unknown | 0 | 0 | 0 |
| Total | 150,962 | 173,837 | 324,799 |

| Antibiotic Utilization | | | | | | | | |
|------------------------|--------------|--------------------------|--------------------------------------|--|--|--|---|--|
| Age | Sex | Total Antibiotic Scripts | Average Scripts PMPY for Antibiotics | Total Days Supplied for All Antibiotic Scripts | Average Days Supplied per Antibiotic Scrip | Total Number of Scripts for Antibiotics of Concern | Average Scripts PMPY for Antibiotics of Concern | Percentage of Antibiotics of Concern of all Antibiotic Scripts |
| 0-9 | M | 6582 | 0.96 | 57957 | 8.81 | 2900 | 0.42 | 44% |
| | F | 6139 | 0.95 | 54203 | 8.83 | 2519 | 0.39 | 41% |
| | Total | 12,721 | 0.95 | 112,160 | 8.82 | 5,419 | 0.41 | 43% |
| 10-17 | M | 1880 | 0.56 | 18789 | 9.99 | 866 | 0.26 | 46% |
| | F | 2480 | 0.76 | 22412 | 9.04 | 1058 | 0.32 | 43% |
| | Total | 4,360 | 0.66 | 41,201 | 9.45 | 1,924 | 0.29 | 44% |
| 18-34 | M | 958 | 0.75 | 9030 | 9.43 | 413 | 0.32 | 43% |
| | F | 4861 | 1.46 | 39759 | 8.18 | 1912 | 0.57 | 39% |
| | Total | 5,819 | 1.26 | 48,789 | 8.38 | 2,325 | 0.51 | 40% |

| | | | | | | | | |
|---------|-------|--------|------|---------|-------|--------|------|-----|
| 35-49 | M | 662 | 1.09 | 7142 | 10.79 | 306 | 0.51 | 46% |
| | F | 1651 | 1.63 | 14365 | 8.70 | 813 | 0.80 | 49% |
| | Total | 2,313 | 1.43 | 21,507 | 9.30 | 1,119 | 0.69 | 48% |
| 50-64 | M | 546 | 1.10 | 5631 | 10.31 | 220 | 0.44 | 40% |
| | F | 617 | 1.56 | 5261 | 8.53 | 280 | 0.71 | 45% |
| | Total | 1,163 | 1.31 | 10,892 | 9.37 | 500 | 0.56 | 43% |
| 65-74 | M | 0 | 0.00 | 0 | NA | 0 | 0.00 | NA |
| | F | 1 | 2.40 | 30 | 30.00 | 0 | 0.00 | 0% |
| | Total | 1 | 1.50 | 30 | 30.00 | 0 | 0.00 | 0% |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | NA |
| Total | M | 10,628 | 0.84 | 98,549 | 9.27 | 4,705 | 0.37 | 44% |
| | F | 15,749 | 1.09 | 136,030 | 8.64 | 6,582 | 0.45 | 42% |
| | Total | 26,377 | 0.97 | 234,579 | 8.89 | 11,287 | 0.42 | 43% |

| Antibiotics of Concern Utilization | | | | | | | | | | | | | | | |
|------------------------------------|--------------|-------------------------|-------------------------------------|--|--|---|--|---------------------------------------|---|-------------------------|------------------------------------|---------------------------|---------------------------------------|--|---|
| Age | Sex | Total Quinolone Scripts | Average Scripts PMPY for Quinolones | Total Cephalosporin 2nd-4th Generation Scripts | Average Scripts PMPY for Cephalosporins 2nd-4th Generation | Total Azithromycin and Clarithromycin Scripts | Average Scripts PMPY for Azithromycins and Clarithromycins | Total Amoxicillin/Clavulanate Scripts | Average Scripts PMPY for Amoxicillin/Clavulanates | Total Ketolides Scripts | Average Scripts PMPY for Ketolides | Total Clindamycin Scripts | Average Scripts PMPY for Clindamycins | Total Misc. Antibiotics of Concern Scripts | Average Scripts PMPY for Misc. Antibiotics of Concern |
| 0-9 | M | 2 | 0.00 | 422 | 0.06 | 1673 | 0.24 | 769 | 0.11 | 0 | 0.00 | 34 | 0.00 | 0 | 0.00 |
| | F | 3 | 0.00 | 350 | 0.05 | 1408 | 0.22 | 708 | 0.11 | 0 | 0.00 | 49 | 0.01 | 1 | 0.00 |
| | Total | 5 | 0.00 | 772 | 0.06 | 3,081 | 0.23 | 1,477 | 0.11 | 0 | 0.00 | 83 | 0.01 | 1 | 0.00 |
| 10-17 | M | 20 | 0.01 | 45 | 0.01 | 561 | 0.17 | 206 | 0.06 | 0 | 0.00 | 29 | 0.01 | 5 | 0.00 |
| | F | 64 | 0.02 | 75 | 0.02 | 674 | 0.21 | 204 | 0.06 | 0 | 0.00 | 39 | 0.01 | 2 | 0.00 |
| | Total | 84 | 0.01 | 120 | 0.02 | 1,235 | 0.19 | 410 | 0.06 | 0 | 0.00 | 68 | 0.01 | 7 | 0.00 |
| 18-34 | M | 59 | 0.05 | 16 | 0.01 | 210 | 0.17 | 80 | 0.06 | 0 | 0.00 | 46 | 0.04 | 2 | 0.00 |
| | F | 362 | 0.11 | 66 | 0.02 | 942 | 0.28 | 330 | 0.10 | 0 | 0.00 | 210 | 0.06 | 2 | 0.00 |
| | Total | 421 | 0.09 | 82 | 0.02 | 1,152 | 0.25 | 410 | 0.09 | 0 | 0.00 | 256 | 0.06 | 4 | 0.00 |
| 35-49 | M | 86 | 0.14 | 16 | 0.03 | 106 | 0.17 | 58 | 0.10 | 0 | 0.00 | 39 | 0.06 | 1 | 0.00 |
| | F | 191 | 0.19 | 26 | 0.03 | 387 | 0.38 | 118 | 0.12 | 0 | 0.00 | 89 | 0.09 | 2 | 0.00 |
| | Total | 277 | 0.17 | 42 | 0.03 | 493 | 0.30 | 176 | 0.11 | 0 | 0.00 | 128 | 0.08 | 3 | 0.00 |
| 50-64 | M | 55 | 0.11 | 5 | 0.01 | 83 | 0.17 | 42 | 0.08 | 0 | 0.00 | 33 | 0.07 | 2 | 0.00 |
| | F | 82 | 0.21 | 3 | 0.01 | 128 | 0.32 | 38 | 0.10 | 0 | 0.00 | 28 | 0.07 | 1 | 0.00 |
| | Total | 137 | 0.15 | 8 | 0.01 | 211 | 0.24 | 80 | 0.09 | 0 | 0.00 | 61 | 0.07 | 3 | 0.00 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 222 | 0.02 | 504 | 0.04 | 2,633 | 0.21 | 1,155 | 0.09 | 0 | 0.00 | 181 | 0.01 | 10 | 0.00 |
| | F | 702 | 0.05 | 520 | 0.04 | 3,539 | 0.24 | 1,398 | 0.10 | 0 | 0.00 | 415 | 0.03 | 8 | 0.00 |
| | Total | 924 | 0.03 | 1,024 | 0.04 | 6,172 | 0.23 | 2,553 | 0.09 | 0 | 0.00 | 596 | 0.02 | 18 | 0.00 |

| All Other Antibiotics Utilization | | | | | | | | | | | | | | | | | |
|-----------------------------------|-------|-------------------------------------|---|-----------------------------|---|---|---|--------------------------|--------------------------------------|--|---|-------------------------|-------------------------------------|---------------------------|---------------------------------------|-------------------------------|---|
| Age | Sex | Total Absorbable Sulfonamide Scrips | Average Scrips PMPY for Absorbable Sulfonamides | Total Aminoglycoside Scrips | Average Scrips PMPY for Aminoglycosides | Total 1st Generation Cephalosporin Scrips | Average Scrips PMPY for 1st Generation Cephalosporins | Total Lincosamide Scrips | Average Scrips PMPY for Lincosamides | Total Macrolides (not azith. or clarith.) Scrips | Average Scrips PMPY for Macrolides (not azith. or clarith.) | Total Penicillin Scrips | Average Scrips PMPY for Penicillins | Total Tetracycline Scrips | Average Scrips PMPY for Tetracyclines | Total Misc. Antibiotic Scrips | Average Scrips PMPY for Misc. Antibiotics |
| 0-9 | M | 321 | 0.05 | 1 | 0.00 | 405 | 0.06 | 0 | 0.00 | 14 | 0.00 | 2919 | 0.43 | 0 | 0.00 | 22 | 0.00 |
| | F | 382 | 0.06 | 0 | 0.00 | 500 | 0.08 | 0 | 0.00 | 6 | 0.00 | 2669 | 0.41 | 7 | 0.00 | 56 | 0.01 |
| | Total | 703 | 0.05 | 1 | 0.00 | 905 | 0.07 | 0 | 0.00 | 20 | 0.00 | 5,588 | 0.42 | 7 | 0.00 | 78 | 0.01 |
| 10-17 | M | 84 | 0.03 | 0 | 0.00 | 194 | 0.06 | 0 | 0.00 | 5 | 0.00 | 566 | 0.17 | 157 | 0.05 | 8 | 0.00 |
| | F | 196 | 0.06 | 3 | 0.00 | 301 | 0.09 | 0 | 0.00 | 11 | 0.00 | 652 | 0.20 | 132 | 0.04 | 127 | 0.04 |
| | Total | 280 | 0.04 | 3 | 0.00 | 495 | 0.07 | 0 | 0.00 | 16 | 0.00 | 1,218 | 0.18 | 289 | 0.04 | 135 | 0.02 |
| 18-34 | M | 85 | 0.07 | 0 | 0.00 | 92 | 0.07 | 0 | 0.00 | 5 | 0.00 | 256 | 0.20 | 88 | 0.07 | 19 | 0.01 |
| | F | 372 | 0.11 | 0 | 0.00 | 502 | 0.15 | 0 | 0.00 | 11 | 0.00 | 920 | 0.28 | 256 | 0.08 | 888 | 0.27 |
| | Total | 457 | 0.10 | 0 | 0.00 | 594 | 0.13 | 0 | 0.00 | 16 | 0.00 | 1,176 | 0.26 | 344 | 0.07 | 907 | 0.20 |
| 35-49 | M | 65 | 0.11 | 0 | 0.00 | 59 | 0.10 | 0 | 0.00 | 6 | 0.01 | 176 | 0.29 | 34 | 0.06 | 16 | 0.03 |
| | F | 144 | 0.14 | 0 | 0.00 | 153 | 0.15 | 0 | 0.00 | 1 | 0.00 | 275 | 0.27 | 91 | 0.09 | 174 | 0.17 |
| | Total | 209 | 0.13 | 0 | 0.00 | 212 | 0.13 | 0 | 0.00 | 7 | 0.00 | 451 | 0.28 | 125 | 0.08 | 190 | 0.12 |
| 50-64 | M | 98 | 0.20 | 0 | 0.00 | 51 | 0.10 | 0 | 0.00 | 0 | 0.00 | 124 | 0.25 | 38 | 0.08 | 15 | 0.03 |
| | F | 75 | 0.19 | 0 | 0.00 | 64 | 0.16 | 0 | 0.00 | 0 | 0.00 | 104 | 0.26 | 29 | 0.07 | 65 | 0.16 |
| | Total | 173 | 0.19 | 0 | 0.00 | 115 | 0.13 | 0 | 0.00 | 0 | 0.00 | 228 | 0.26 | 67 | 0.08 | 80 | 0.09 |
| 65-74 | M | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 |
| | F | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 2.40 |
| | Total | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 0 | 0.00 | 1 | 1.50 |
| 75-84 | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| 85+ | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Unknown | M | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | F | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| | Total | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA | 0 | NA |
| Total | M | 653 | 0.05 | 1 | 0.00 | 801 | 0.06 | 0 | 0.00 | 30 | 0.00 | 4,041 | 0.32 | 317 | 0.03 | 80 | 0.01 |
| | F | 1,169 | 0.08 | 3 | 0.00 | 1,520 | 0.10 | 0 | 0.00 | 29 | 0.00 | 4,620 | 0.32 | 515 | 0.04 | 1,311 | 0.09 |
| | Total | 1,822 | 0.07 | 4 | 0.00 | 2,321 | 0.09 | 0 | 0.00 | 59 | 0.00 | 8,661 | 0.32 | 832 | 0.03 | 1,391 | 0.05 |

| Board Certification (BCR) | | | |
|---|------------------------------|---------------------|---------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Type of Physician | Number of Physicians in Each | Board Certification | |
| | | Number | Percent |
| Family Medicine | NR | NR | NR |
| Internal Medicine | NR | NR | NR |
| OB/GYN physicians | NR | NR | NR |
| Pediatricians | NR | NR | NR |
| Geriatricians | NR | NR | NR |
| Other physician specialists | NR | NR | NR |

| Total Membership (TLM) | |
|--|---------------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| Product/Product Line | Total Number of Members* |
| HMO (Total) | 30,164 |
| Medicaid | 30164 |
| Commercial | 0 |
| Medicare (cost or risk) | 0 |
| Other | 0 |
| PPO (Total) | 0 |
| Medicaid | 0 |
| Commercial | 0 |
| Medicare (cost or risk) | 0 |
| Other | 0 |
| POS (Total) | 0 |
| Medicaid | 0 |
| Commercial | 0 |
| Medicare (cost or risk) | 0 |
| Other | 0 |
| FFS (Total) | 0 |
| Medicaid | 0 |
| Commercial | 0 |
| Medicare (cost or risk) | 0 |
| Other | 0 |
| Total | 30,164 |
| * Total number of members in each category as of December 31 of the measurement year. | |

| Enrollment by Product Line: Total (ENPA) | | | |
|---|---------------------------|-----------------------------|----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Age | Male Member Months | Female Member Months | Total Member Months |
| <1 | 12301 | 11580 | 23,881 |
| 1-4 | 37578 | 36415 | 73,993 |
| 5-9 | 32302 | 29796 | 62,098 |
| 10-14 | 26343 | 25598 | 51,941 |
| 15-17 | 13966 | 13582 | 27,548 |
| 18-19 | 4661 | 6036 | 10,697 |
| 0-19 Subtotal | 127,151 | 123,007 | 250,158 |
| 0-19 Subtotal: % | 84.23% | 70.76% | 77.02% |
| 20-24 | 3103 | 12860 | 15,963 |
| 25-29 | 3694 | 12146 | 15,840 |
| 30-34 | 3802 | 8934 | 12,736 |
| 35-39 | 2738 | 6014 | 8,752 |
| 40-44 | 2284 | 3638 | 5,922 |
| 20-44 Subtotal | 15,621 | 43,592 | 59,213 |
| 20-44 Subtotal: % | 10.35% | 25.08% | 18.23% |
| 45-49 | 2248 | 2492 | 4,740 |
| 50-54 | 2268 | 2066 | 4,334 |
| 55-59 | 2154 | 1535 | 3,689 |
| 60-64 | 1517 | 1140 | 2,657 |
| 45-64 Subtotal | 8,187 | 7,233 | 15,420 |
| 45-64 Subtotal: % | 5.42% | 4.16% | 4.75% |
| 65-69 | 2 | 1 | 3 |
| 70-74 | 1 | 4 | 5 |
| 75-79 | 0 | 0 | 0 |
| 80-84 | 0 | 0 | 0 |
| 85-89 | 0 | 0 | 0 |
| >=90 | 0 | 0 | 0 |
| >=65 Subtotal | 3 | 5 | 8 |
| >=65 Subtotal: % | 0.00% | 0.00% | 0.00% |
| Age Unknown | 0 | 0 | 0 |
| Total | 150,962 | 173,837 | 324,799 |

| Enrollment by Product Line: Dual Eligibles (ENPB) | | | |
|--|--------------------|----------------------|---------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Age | Male Member Months | Female Member Months | Total Member Months |
| <1 | NR | NR | NR |
| 1-4 | NR | NR | NR |
| 5-9 | NR | NR | NR |
| 10-14 | NR | NR | NR |
| 15-17 | NR | NR | NR |
| 18-19 | NR | NR | NR |
| 0-19 Subtotal | NR | NR | NR |
| 0-19 Subtotal: % | NR | NR | NR |
| 20-24 | NR | NR | NR |
| 25-29 | NR | NR | NR |
| 30-34 | NR | NR | NR |
| 35-39 | NR | NR | NR |
| 40-44 | NR | NR | NR |
| 20-44 Subtotal | NR | NR | NR |
| 20-44 Subtotal: % | NR | NR | NR |
| 45-49 | NR | NR | NR |
| 50-54 | NR | NR | NR |
| 55-59 | NR | NR | NR |
| 60-64 | NR | NR | NR |
| 45-64 Subtotal | NR | NR | NR |
| 45-64 Subtotal: % | NR | NR | NR |
| 65-69 | NR | NR | NR |
| 70-74 | NR | NR | NR |
| 75-79 | NR | NR | NR |
| 80-84 | NR | NR | NR |
| 85-89 | NR | NR | NR |
| >=90 | NR | NR | NR |
| >=65 Subtotal | NR | NR | NR |
| >=65 Subtotal: % | NR | NR | NR |
| Age Unknown | NR | NR | NR |
| Total | NR | NR | NR |

Enrollment by Product Line: Disabled (ENPC)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Age | Male Member Months | Female Member Months | Total Member Months |
|-------------------|--------------------|----------------------|---------------------|
| <1 | NR | NR | NR |
| 1-4 | NR | NR | NR |
| 5-9 | NR | NR | NR |
| 10-14 | NR | NR | NR |
| 15-17 | NR | NR | NR |
| 18-19 | NR | NR | NR |
| 0-19 Subtotal | NR | NR | NR |
| 0-19 Subtotal: % | NR | NR | NR |
| 20-24 | NR | NR | NR |
| 25-29 | NR | NR | NR |
| 30-34 | NR | NR | NR |
| 35-39 | NR | NR | NR |
| 40-44 | NR | NR | NR |
| 20-44 Subtotal | NR | NR | NR |
| 20-44 Subtotal: % | NR | NR | NR |
| 45-49 | NR | NR | NR |
| 50-54 | NR | NR | NR |
| 55-59 | NR | NR | NR |
| 60-64 | NR | NR | NR |
| 45-64 Subtotal | NR | NR | NR |
| 45-64 Subtotal: % | NR | NR | NR |
| 65-69 | NR | NR | NR |
| 70-74 | NR | NR | NR |
| 75-79 | NR | NR | NR |
| 80-84 | NR | NR | NR |
| 85-89 | NR | NR | NR |
| >=90 | NR | NR | NR |
| >=65 Subtotal | NR | NR | NR |
| >=65 Subtotal: % | NR | NR | NR |
| Age Unknown | NR | NR | NR |
| Total | NR | NR | NR |

| Enrollment by Product Line: Other (ENPD) | | | |
|--|---------------------------|-----------------------------|----------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Age | Male Member Months | Female Member Months | Total Member Months |
| <1 | NR | NR | NR |
| 1-4 | NR | NR | NR |
| 5-9 | NR | NR | NR |
| 10-14 | NR | NR | NR |
| 15-17 | NR | NR | NR |
| 18-19 | NR | NR | NR |
| 0-19 Subtotal | NR | NR | NR |
| 0-19 Subtotal: % | NR | NR | NR |
| 20-24 | NR | NR | NR |
| 25-29 | NR | NR | NR |
| 30-34 | NR | NR | NR |
| 35-39 | NR | NR | NR |
| 40-44 | NR | NR | NR |
| 20-44 Subtotal | NR | NR | NR |
| 20-44 Subtotal: % | NR | NR | NR |
| 45-49 | NR | NR | NR |
| 50-54 | NR | NR | NR |
| 55-59 | NR | NR | NR |
| 60-64 | NR | NR | NR |
| 45-64 Subtotal | NR | NR | NR |
| 45-64 Subtotal: % | NR | NR | NR |
| 65-69 | NR | NR | NR |
| 70-74 | NR | NR | NR |
| 75-79 | NR | NR | NR |
| 80-84 | NR | NR | NR |
| 85-89 | NR | NR | NR |
| >=90 | NR | NR | NR |
| >=65 Subtotal | NR | NR | NR |
| >=65 Subtotal: % | NR | NR | NR |
| Age Unknown | NR | NR | NR |
| Total | NR | NR | NR |

| Enrollment by State (EBS) | |
|--|---------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | |
| State | Number |
| Alabama | 0 |
| Alaska | 0 |
| Arizona | 37 |
| Arkansas | 0 |
| California | 3 |
| Colorado | 12 |
| Connecticut | 0 |
| Delaware | 0 |
| District of Columbia | 0 |
| Florida | 0 |
| Georgia | 0 |
| Hawaii | 0 |
| Idaho | 0 |
| Illinois | 1 |
| Indiana | 0 |
| Iowa | 2 |
| Kansas | 0 |
| Kentucky | 1 |
| Louisiana | 0 |
| Maine | 0 |
| Maryland | 0 |
| Massachusetts | 1 |
| Michigan | 0 |
| Minnesota | 0 |
| Mississippi | 0 |
| Missouri | 0 |
| Montana | 0 |
| Nebraska | 0 |
| Nevada | 1 |
| New Hampshire | 1 |
| New Jersey | 0 |
| New Mexico | 30080 |
| New York | 0 |
| North Carolina | 1 |
| North Dakota | 0 |
| Ohio | 0 |
| Oklahoma | 2 |
| Oregon | 0 |
| Pennsylvania | 0 |
| Rhode Island | 0 |
| South Carolina | 0 |
| South Dakota | 1 |
| Tennessee | 0 |
| Texas | 20 |
| Utah | 0 |
| Vermont | 0 |
| Virginia | 1 |
| Washington | 0 |
| West Virginia | 0 |
| Wisconsin | 0 |
| Wyoming | 0 |
| American Samoa | 0 |
| Federated States of Micronesia | 0 |
| Guam | 0 |
| Commonwealth of Northern Marianas | 0 |
| Puerto Rico | 0 |
| Virgin Islands | 0 |
| Other | 0 |
| TOTAL | 30,164 |

Race/Ethnicity Diversity of Membership (RDM)

Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None)

| Race/Ethnicity Diversity of Membership | |
|---|----|
| Total Unduplicated Membership During the Measurement Year | NR |

Percentage of Members for Whom the Organization has Race/Ethnicity Information by Data Collection

| Direct Data Collection Method | | | Indirect Data Collection Method | | Unknown | |
|-------------------------------|---------------------|----|---------------------------------|----|---------|----|
| Race | Direct Total | NR | Indirect Total* | NR | Total* | NR |
| | Health Plan Direct* | NR | | | | |
| | CMS/State Database* | NR | | | | |
| | Other* | NR | | | | |
| Ethnicity | Direct Total | NR | Indirect Total* | NR | Total* | NR |
| | Health Plan Direct* | NR | | | | |
| | CMS/State Database* | NR | | | | |
| | Other* | NR | | | | |

*Enter percentage as a value between 0 and 1.

| Race | Hispanic or Latino | | Not Hispanic or Latino | | Unknown Ethnicity | | Declined Ethnicity | | Total | |
|---|--------------------|------------|------------------------|------------|-------------------|------------|--------------------|------------|--------|------------|
| | Number | Percentage | Number | Percentage | Number | Percentage | Number | Percentage | Number | Percentage |
| White | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Black or African American | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| American-Indian and Alaska Native | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Asian | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Native Hawaiian and Other Pacific Islanders | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Some Other Race | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Two or More Races | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Unknown | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Declined | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Total | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

| Language Diversity of Membership (LDM) | | | |
|--|---------------------------|----------------------------|---------------------------------|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | | |
| Percentage of Members With Known Language Value from Each Data Source | | | |
| Category | Health Plan Direct | CMS/State Databases | Other Third-Party Source |
| Spoken Language Preferred for Health Care* | NR | NR | NR |
| Preferred Language for Written Materials* | NR | NR | NR |
| Other Language Needs* | NR | NR | NR |
| *Enter percentage as a value between 0 and 1. | | | |
| Spoken Language Preferred for Health Care | | | |
| | Number | Percentage | |
| English | NR | NR | |
| Non-English | NR | NR | |
| Unknown | NR | NR | |
| Declined | NR | NR | |
| Total* | NR | NR | |
| Language Preferred for Written Materials | | | |
| | Number | Percentage | |
| English | NR | NR | |
| Non-English | NR | NR | |
| Unknown | NR | NR | |
| Declined | NR | NR | |
| Total* | NR | NR | |
| Other Language Needs | | | |
| | Number | Percentage | |
| English | NR | NR | |
| Non-English | NR | NR | |
| Unknown | NR | NR | |
| Declined | NR | NR | |
| Total* | NR | NR | |
| *Should sum to 100% | | | |

| | | |
|--|--|--|
| Weeks of Pregnancy at Time of Enrollment in MCO (WOP) | | |
|--|--|--|

| | | |
|--|--|--|
| Blue Cross and Blue Shield of New Mexico (Org ID: 1825, SubID: 9522, Medicaid, Spec Area: None, Spec Proj: None) | | |
|--|--|--|

| Measurement Year | | |
|--------------------|--------------|----------------|
| Measurement Year | 2012 | |
| Weeks of Pregnancy | Number | Percentage |
| < 0 weeks | 299 | 21.79% |
| 1-12 weeks | 117 | 8.53% |
| 13-27 weeks | 616 | 44.90% |
| 28 or more weeks | 250 | 18.22% |
| Unknown | 90 | 6.56% |
| Total | 1,372 | 100.00% |